

# Cosmology: Example Sheet 3

David Tong, November 2019

1. A non-relativistic, quantum mechanical particle sits in a box whose sides have length  $a(t) \times L$ . Write down the wavefunctions when  $a(t)$  is constant.

It can be shown that if  $a(t)$  changes suitably slowly then the system remains in a given energy eigenstate. Show that the momentum “redshifts” as  $\mathbf{p}(t) = \mathbf{p}_0/a(t)$  where  $\mathbf{p}_0$  is the momentum when  $a(t_0) = 1$ .

A gas of non-relativistic particles at temperature  $T$  is described by the Maxwell-Boltzmann distribution at time  $t = t_0$ . Assuming the momentum-redshift above, show that the gas retains the Maxwell-Boltzmann form as the universe expands, but with a temperature that scales as  $T(t) = T_0/a(t)^2$ .

2. The thermal cosmic microwave background is assumed to be isotropic with a temperature  $T$  in an inertial frame  $S$ . The same radiation is detected in another inertial frame  $S'$ , moving with velocity  $v$  with respect to  $S$ .

The Lorentz transformation relating the energy  $E$  and 3-momentum  $\mathbf{p}$  of a particle in the two frames is

$$E = \gamma(E' - \mathbf{v} \cdot \mathbf{p}')$$

where  $\gamma = 1/\sqrt{1 - v^2/c^2}$ . A photon has  $E = pc$ . Show that the microwave background will also appear thermal in  $S'$ , but with an anisotropic temperature

$$T'(\theta') = \frac{T}{\gamma(1 - (v/c) \cos \theta')} = T \left( 1 + \frac{v}{c} \cos \theta' + \mathcal{O}(v^2/c^2) \right)$$

where  $\theta'$  is the angle between the velocity  $\mathbf{v}$  and the momentum  $\mathbf{p}'$  of the photon arriving at the detector.

Let  $T'_+$  and  $T'_-$  be the maximum and minimum temperatures seen in the inertial frame  $S'$ . Show that

$$T = \sqrt{T'_+ T'_-}$$

The observed CMB, shown in the figure, has  $T'_+ - T'_- \approx 6.5 \times 10^{-3}$  K, with  $T = \sqrt{T'_+ T'_-} \approx 2.7$  K. How fast are we travelling with respect to the universe’s preferred inertial frame?




Figure 1: The observed CMB dipole.

It is believed that there exists a yet-to-be-observed thermal cosmic neutrino background that is isotropic in the same frame  $S$  as the CMB. The neutrino has a small mass and so  $E^2 = p^2c^2 + m^2c^4$ . Today, the neutrinos are travelling at non-relativistic speeds. Show that when (if!) we finally observe the cosmic neutrino background, we do not expect the energy density to be thermal, even at a fixed angle.

**3.** The Planck blackbody formula states that the number of photons with frequency between  $\omega$  and  $\omega + d\omega$  is

$$n(\omega) d\omega = \frac{1}{\pi^2 c^3} \frac{\omega^2}{e^{\beta\hbar\omega} - 1}$$

Show that the total number of photons is

$$n_\gamma = \frac{2\zeta(3)}{\pi^2 \hbar^3 c^3} (k_B T)^3 \quad (1)$$

[You will need the integral  $\int_0^\infty dy y^2/(e^y - 1) = 2\zeta(3)$  with  $\zeta(3) \approx 1.3$ .]

Define  $n_{\text{energetic}}$  to be the number of energetic photons with energy greater than the hydrogen binding  $E_{\text{bind}} \approx 13.6$  eV. Show that, when  $k_B T \ll E_{\text{bind}}$ ,

$$\frac{n_{\text{energetic}}}{n_\gamma} \approx \frac{(\beta E_{\text{bind}})^2}{2\zeta(3)} e^{-\beta E_{\text{bind}}}$$

As a naive diagnostic, suppose that recombination occurs when there is less than a single energetic photon per baryon. Use the baryon-to-photon ratio  $\eta = n_B/n_\gamma \approx 10^{-9}$  to determine the temperature and redshift of recombination according to this criterion.

4. Assume that electrons, protons and hydrogen are in chemical equilibrium during recombination, with the chemical potentials related by  $\mu_e + \mu_p = \mu_H$ . Show that the number of electrons is related to the number of hydrogen atoms by

$$n_e^2 \approx n_H \left( \frac{m_e k_B T}{2\pi \hbar^2} \right)^{3/2} e^{-\beta E_{\text{bind}}}$$

with  $E_{\text{bind}}$  the binding energy of the hydrogen ground state. What assumptions did you make along the way?

The ionization fraction is defined as  $X_e = n_e/n_B$  with  $n_B \approx n_p + n_H$ , the number of baryons. Use the expression (1) to show that

$$\frac{1 - X_e}{X_e^2} = \eta \frac{2\zeta(3)}{\pi^2} \left( \frac{2\pi k_B T}{m_e c^2} \right)^{3/2} e^{\beta E_{\text{bind}}}$$

with  $\eta$  the baryon-to-photon ratio. Consider the limiting regimes  $k_B T \ll E_{\text{bind}}$  and  $E_{\text{bind}} \ll k_B T \ll m_e c^2$  to roughly sketch  $X_e$  as a function of temperature.

5. Recombination is not instantaneous, but happens over a period of time. Some photons in the CMB come from earlier times, when the universe was hotter, and some from later times.

Why does the observed CMB exhibit a perfect blackbody spectrum at a single temperature?

6. At temperature  $T$ , and vanishing chemical potential, the expected number of particles with momentum  $\mathbf{p}$  is given by

$$n(\mathbf{p}) = \frac{1}{e^{\beta E(\mathbf{p})} \mp 1}$$

where the minus sign is for bosons and the plus sign for fermions.

For ultra-relativistic particles, with  $E(\mathbf{p}) = pc$ , show that the total number of fermions,  $n_F$ , is related to the total number of bosons,  $n_B$ , by  $n_F = 3n_B/4$ . Show that the total energy density of fermions,  $\rho_F$ , is related to the total energy density of bosons,  $\rho_B$ , by  $\rho_F = 7\rho_B/8$ .

[Note: you need not evaluate any integral to do this question.]

7. Consider a gas of electrons and positrons in the ultra-relativistic limit  $k_B T \gg m_e c^2$ . In the early universe, there must have been a slight imbalance of electrons over positrons. This is modelled by introducing a small chemical potential  $\mu_e \ll k_B T$  for electrons, with an equal and opposite chemical potential for positrons. Show that this results in a small excess  $\Delta n$  of electrons over positrons, given by

$$\Delta n = \frac{g(k_B T)^2 \mu_e}{6\hbar^3 c^3} \left( 1 + \mathcal{O} \left( \frac{\mu_e^2}{k_B^2 T^2} \right) \right)$$

[You will need to use the integral  $\int_0^\infty dy y/(e^y + 1) = \pi^2/12$ .]