

# The Standard Model: Example Sheet 3

David Tong, February 2026

1. Show that there is no consistent *chiral*  $U(1)$  gauge theory with  $N = 4$  Weyl fermions with integer charges that can be coupled to gravity.

Find an example of a consistent  $U(1)$  gauge theory with  $N = 5$  Weyl fermions with integer charges that can be coupled to gravity.

2\*. An  $SU(N)$  gauge theory is coupled to a single left-handed Weyl fermion  $\chi$  in the symmetric  $\square\square$  representation and  $p$  left-handed Weyl fermions  $\psi_i$ ,  $i = 1, \dots, p$ , in the anti-fundamental  $\bar{\square}$  representation. For what value of  $p$  is the quantum theory consistent?

Classically, the theory has a  $G_F = SU(p)$  global symmetry that acts on the  $\psi_i$ . In addition, there are two  $U(1)$  symmetries

$$\begin{aligned} U(1)_\chi : \quad & \chi \rightarrow e^{i\alpha} \chi \quad \text{and} \quad \psi_i \rightarrow \psi_i \\ U(1)_\psi : \quad & \chi \rightarrow \chi \quad \text{and} \quad \psi_i \rightarrow e^{i\beta} \psi_i \end{aligned}$$

Show that each of the  $U(1)$  symmetries suffers a chiral anomaly and so is not a symmetry of the quantum theory. Find a linear combination of these symmetries that does survive in the quantum theory.

Compute the  $SU(p)^3$ ,  $SU(p)^2 U(1)$ ,  $U(1)^3$  and  $U(1)$  (i.e. mixed gauge-gravity) 't Hooft anomalies for the symmetries of the quantum theory.

It is conjectured that this theory confines without spontaneously breaking any global symmetry. The massless fields are thought to be a collection of gauge singlet fermions

$$\lambda^{ij} = \psi^{[i} (\chi \psi^{j]})$$

transforming in the anti-symmetric  $\square\square$  representation of the  $SU(p)$  global symmetry. What is the  $U(1)$  charge of this fermion? Show that the 't Hooft anomalies of the fermion  $\lambda$  match those of the original gauge theory.

Note: You may find the following data helpful. For  $SU(N)$ , the fundamental  $\square$ , adjoint, symmetric  $\square\square$  and anti-symmetric  $\square\square$  have the dimension, Dynkin index  $I(R) = I(\bar{R})$  and anomaly coefficient  $A(R) = -A(\bar{R})$  given by

$R$	$\square$	adj	$\square\square$	$\square\square$
$\dim(R)$	$N$	$N^2 - 1$	$\frac{1}{2}N(N + 1)$	$\frac{1}{2}N(N - 1)$
$I(R)$	1	$2N$	$N + 2$	$N - 2$
$A(R)$	1	0	$N + 4$	$N - 4$

**3\***. Consider one generation of the Standard Model, with left- and right-handed Weyl fermions transforming in the following representations of  $U(1) \times SU(2) \times SU(3)$ :

$$Q_L : (\mathbf{2}, \mathbf{3})_q, \quad L_L : (\mathbf{2}, \mathbf{1})_l, \quad u_R : (\mathbf{1}, \mathbf{3})_u, \quad d_R : (\mathbf{1}, \mathbf{3})_d, \quad e_R : (\mathbf{1}, \mathbf{1})_x .$$

This is the usual set of representations (ignoring the right-handed neutrino), but with the hypercharges  $q, l, u, d$ , and  $x$  left arbitrary. Assume that all hypercharges are real numbers, i.e.  $q, l, u, d, x \in \mathbb{R}$

i) Write down all the conditions for anomaly cancellation, including the mixed  $U(1)$ -gravitational anomaly.

ii) Show that these equations have two solutions, one with  $u = -d$  and the other the hypercharges of the Standard Model (up to scaling).

**4.** This is a repeat of question 4, but with the additional assumption that hypercharges are integers, i.e.  $q, l, u, d, x \in \mathbb{Z}$ .

i) Write down all conditions for anomaly cancellation, this time omitting the requirement that the theory can be coupled to gravity.

ii) Show that there is a unique solution to these equations that automatically satisfies the mixed gauge-gravitational anomaly. (You may invoke, without proof, any pure mathematics result that your grandmother has heard of.)

[Hint: First argue that you can write  $u - d = 2y$  for  $y \in \mathbb{Z}$ . Then use the change of variables

$$x = -\frac{6}{v+w} \quad \text{and} \quad y = \frac{3(v-w)}{v+w} . ]$$

**5.** What are the global symmetries of the classical Lagrangian of the Standard Model assuming:

i) there are no right-handed neutrinos

ii) right-handed neutrinos exist, with all possible relevant and marginal interactions included?

How do these conclusions change if you include the effect of anomalies?

**6a.** [This question is optional. I didn't discuss surviving discrete symmetries in class this year because it's something of a distraction from our main interest, namely the Standard Model. Still, this is a nice question that you might try if you are interested in more formal aspects of physics.]

Consider  $SU(N)$  Yang-Mills coupled to a single, massless left-handed Weyl fermion  $\lambda$  in the adjoint representation. Why does this make sense as a quantum theory? (As an aside: this theory happens to enjoy  $\mathcal{N} = 1$  supersymmetry, although this fact is not needed for this question.)

Classically this theory has a  $U(1)$  global symmetry which acts as  $\lambda \rightarrow e^{i\alpha}\lambda$ . By considering how this transformation affects the  $\theta$  angle, show that a  $\mathbb{Z}_{2N}$  subgroup survives in the quantum theory.

The theory is expected to confine, develop a mass gap, and form a condensate

$$\langle \text{Tr } \lambda\lambda \rangle \sim \Lambda_{\text{QCD}}^3 . \quad (1)$$

How does the  $\mathbb{Z}_{2N}$  global symmetry act on this condensate? How many ground states does the theory have?

**b.** Consider  $SU(N)$  Yang-Mills (with  $N > 2$ ) coupled to a single *Dirac* fermion  $\psi$  in the symmetric  $\square\square$  representation. What are the classical global symmetries? What are the quantum global symmetries?

This theory is expected to confine, develop a mass gap, and form a condensate

$$\langle \bar{\psi}\psi \rangle \sim \Lambda_{\text{QCD}}^3$$

How many ground states does the theory have? How do your answers change if the Dirac fermion is in the anti-symmetric  $\square$  representation of  $SU(N)$ ?

[Note: You may need to refer to the table in Question 2.]