

# Supersymmetric Quantum Mechanics

---

**David Tong**

*Department of Applied Mathematics and Theoretical Physics,  
Centre for Mathematical Sciences,  
Wilberforce Road,  
Cambridge, CB3 0BA, UK*

<http://www.damtp.cam.ac.uk/user/tong/susyqm.html>

d.tong@damtp.cam.ac.uk

## Recommended Books and Resources

There are very few decent textbooks that cover the material of these lectures. The handful of textbooks that exist with titles like “supersymmetric quantum mechanics” tend to focus on the slightly dull topics of exact solutions, rather than on the connections to geometry that we care about here. Nonetheless, there are two books that will be useful for what lies ahead:

- Nakahara “*Geometry, Topology and Physics*”

This book covers homology, cohomology, index theorems and Kähler manifolds, which is much of the mathematics you’ll need in these lectures. Later editions of the book also cover supersymmetric quantum mechanics towards the end although, in contrast to the rest of the book, the presentation of this material isn’t particularly good.

- Kentaro Hori, in the Clay Mathematics Monograph “*Mirror Symmetry*”.

This book is something of a mixed bag, with contributions from many authors. But the sections written by Kentaro Hori, which comprise Part 2 and Part 3 of the book (pages 143 to 480) are spectacularly good. Our lectures will largely follow the first few steps along the path laid down by Kentaro although I suspect that he would disapprove of the times I replace his rigorous statements with wild, but enthusiastic, handwaving. You can download the book directly from the [Clay Mathematics Institute](#).

While decent books on the subject are in short supply, there is one resource that I strongly recommend. A remarkably large fraction of these lectures (not to mention subsequent developments in the field) is due to Edward Witten. His papers are not only brimming with beautiful physics, but are also models of scientific writing. If you want to learn large swathes of modern physics, you could do worse than turn to Witten’s papers. Those from the late 1970s and early 1980s are particularly accessible. Much of what we cover in these lectures can be found in the papers “[Constraints on Supersymmetry Breaking](#)” and “[Supersymmetry and Morse Theory](#)”. These, and a number of further resources, can be found on the [course webpage](#).

---

# Contents

<b>0</b>	<b>Introduction</b>	<b>3</b>
<b>1</b>	<b>Introducing Supersymmetric Quantum Mechanics</b>	<b>5</b>
1.1	Supersymmetry Algebra	5
1.1.1	A First Look at the Energy Spectrum	5
1.2	A Particle in a Potential	7
1.2.1	Ground States	9
1.2.2	The Witten Index	14
1.3	The Supersymmetric Action	16
1.3.1	Supersymmetry as a Fermionic Symmetry	19
1.4	A Particle Moving in Higher Dimensions	21
1.4.1	A First Look at Morse Theory	21
1.4.2	More Supersymmetry and Holomorphy	25
1.4.3	Less Supersymmetry and Spinors	29
<b>2</b>	<b>Supersymmetry and the Path Integral</b>	<b>33</b>
2.1	The Partition Function and the Index	33
2.1.1	An Example: The Harmonic Oscillator	35
2.1.2	Fermions: Periodic or Anti-Periodic?	38
2.1.3	The Witten Index Revisited	42
2.2	Instantons	46
2.2.1	Tunnelling	49
2.2.2	The Dilute Gas Approximation	54
2.3	Instantons and Supersymmetry	56
2.3.1	Fermi Zero Modes	57
2.3.2	Computing Determinants	59
2.3.3	Computing the Ground State Energy	61
2.3.4	One Last Example: A Particle on a Circle	63
<b>3</b>	<b>Supersymmetry and Geometry</b>	<b>65</b>
3.1	The Supersymmetric Sigma Model	68
3.1.1	Quantisation: Filling in Forms	71
3.1.2	Ground States and de Rham Cohomology	74
3.1.3	The Witten Index and the Chern-Gauss-Bonnet Theorem	80
3.2	Morse Theory	83

3.2.1	Instantons Again	87
3.2.2	The Morse-Witten Complex	95
3.3	The Atiyah-Singer Index Theorem	97
3.3.1	The $N = 1$ Sigma Model	98
3.3.2	The Path Integral Again	100
3.3.3	Adding a Gauge Field	104
3.4	What Comes Next?	106

---

## Acknowledgements

This course is aimed at beginning graduate students. It assumes a good background in quantum mechanics and path integrals and a basic knowledge of differential geometry, say at the level of my lecture notes on [General Relativity](#). The course is based largely on lectures by Kentaro Hori and David Skinner. I'm particularly grateful to David Skinner and Oscar Randal-Williams for patiently explaining a number of issues to me. I also thank Andy Zhao for helping me get minus signs right in the susy variation of the sigma model action.

## 0 Introduction

It will come as no surprise to hear that there is a close relationship between mathematics and physics. Yet, for many centuries, the relationship was more than a little one sided. There was, in the language of marriage counsellors, a lack of equitable reciprocity. Physicists took, but gave little in return. Admittedly there were exceptions, some of them rather important like Newton's development of calculus. Nonetheless, it remains true that mathematics is a tool that us physicists cannot live without, while many mathematicians have no more use of physics than they do of chemistry or botany.

In the last few decades, this narrative has started to change. Physicists have been giving back. As our understanding of quantum field theories has grown, we have uncovered increasingly sophisticated mathematical structures lurking within. These are largely, but not exclusively, the structures that arise in geometry and topology. Using physicist's methods and techniques to solve quantum fields theories has revealed connections to these mathematical ideas. Initially this gave new ways of deriving results well known to mathematicians. But, as the quantum field theories became more involved, so too did the mathematics until physicists were able to discover new results that came as a complete surprise to mathematicians. Prominent among these is an idea called *mirror symmetry*, a novel relationship between different manifolds.

You might reasonably wonder what advantage physicists have over mathematicians in this game. After all, we're certainly not smarter. (At least, not most of us.) And yet, there are times when we are able to leapfrog mathematicians and then turn around and present them with new results that sit firmly within their area of expertise. This seems unfair, like physicists have some kind of secret weapon that mathematicians are unable to wield. And we do. In fact, we have two. The first is the path integral. The second, a wilful disregard for rigour.

These two weapons are not unrelated. The path integral approach to quantum field theory has so far evaded attempts to be placed on a rigorous footing, at least beyond quantum mechanics. This means that most often the physicist's approach to these questions does not meet the mathematician's bar for proof. Physics is perhaps better thought of as an idea generating machine, giving new insights into areas of mathematics that can subsequently be proven using more traditional methods. Happily, in most cases, these subsequent proofs have turned out to be much more than an exercise in dotting i's and crossing  $\hbar$ 's. Mathematicians take their own path to a problem, developing new ideas along the way, and these then feed back into our understanding of quantum field theory. Over the past few decades this process has resulted in a

harmonious and extraordinarily fruitful relationship between communities of physicists and mathematicians.

This interaction has revolutionised certain areas of mathematics. For example, it's difficult to envisage a thriving field of symplectic geometry without mirror symmetry. But it has also changed what we mean by “mathematical physics”. Towards the end of the 20<sup>th</sup> century, this was viewed as a rather dry subject and mostly involved bringing a mathematician's level of pedantry to bear on problems that physicists care about, but with little insight flowing back into the underlying physics. Now, this situation has been reversed, with interesting and exciting ideas flowing in both directions. To emphasise the shift of focus, this new activity is sometimes rebranded “physical mathematics”.

Much of this interplay between physics and mathematics takes place in the arena of supersymmetric field theories. (There are important exceptions, Witten's Fields medal winning work on knot polynomials in Chern Simons theory among them.) Supersymmetric theories are a class of quantum field theories that have a symmetry relating bosons and fermions. There is, so far, no experimental evidence that supersymmetry is a symmetry of our world. But supersymmetric theories have a number of special properties that allow us to make much more progress in solving them than would otherwise be possible. It is often in these solutions to supersymmetric field theories that we find results of interest to mathematicians.

The purpose of these lectures is to take the first few steps along this journey. Sadly we will not reach the heights of the subject like mirror symmetry or knot invariants, both of which require quantum field theories in higher dimensions ( $d = 1 + 1$  and  $d = 2 + 1$  respectively). Instead, we will restrict ourselves to  $d = 0 + 1$  dimensional quantum field theories, also known as quantum mechanics. We will study a number of examples of supersymmetric quantum mechanics and, in solving them, recover some of the highlights of 20<sup>th</sup> century geometry, including ideas of de Rham, Hodge, Morse, Atiyah and Singer.

I should warn you that the level of rigour when addressing the more mathematical aspect of these lectures will be mediocre at best. Anyone with a real interest in these ideas is encouraged to learn both the underlying mathematics and physics to truly appreciate how the two connect. But that is not the path we will take here. Instead, these lectures will assume only a basic knowledge in differential geometry (at the level, say, of my lectures on [General Relativity](#).) We will then use supersymmetric quantum mechanics as a vehicle to take us deeper into the mathematician's territory, allowing us to take a peek at some of the beautiful vistas that await.

# 1 Introducing Supersymmetric Quantum Mechanics

In this section, we discuss some basic facts about supersymmetric quantum mechanics. Our focus will be on a simple class of quantum mechanical systems that, while they have a certain elegance, won't exhibit any deep mathematics. Instead, we will treat them as a proving ground, allowing us to build some intuition for supersymmetry while developing a number of useful calculational techniques. We'll then bring these to bear on problems with a deeper mathematical pedigree in Section 3.

## 1.1 Supersymmetry Algebra

Supersymmetric quantum mechanics is the name given to a class of Hamiltonians  $H$  that can be written as

$$H = \frac{1}{2}\{Q, Q^\dagger\} \quad \text{with} \quad Q^2 = 0 \quad (1.1)$$

Here  $\{A, B\} = AB + BA$  is the anti-commutator. The operator  $Q$  is called the *supercharge* and, as you can see, is something like the square root of the Hamiltonian. Equation (1.1) is called the *supersymmetry algebra*. As we will see, Hamiltonians that can be written in this way enjoy many special properties.

### 1.1.1 A First Look at the Energy Spectrum

The first property is straightforward: the energy of any state is necessarily non-negative. To see this, we just take the usual expectation value in a state  $|\psi\rangle$ ,

$$\begin{aligned} 2\langle\psi|H|\psi\rangle &= \langle\psi|Q^\dagger Q + Q Q^\dagger|\psi\rangle \\ &= |Q|\psi\rangle|^2 + |Q^\dagger|\psi\rangle|^2 \geq 0 \end{aligned}$$

Furthermore, we see that energy  $E$  is only zero for states  $|\psi\rangle$  that are annihilated by both the supercharge and its adjoint

$$E = 0 \quad \Leftrightarrow \quad Q|\psi\rangle = Q^\dagger|\psi\rangle = 0 \quad (1.2)$$

Already, the statement that we have a positive definite spectrum is slightly surprising. Usually in quantum mechanics, we don't care about the overall energy of states since we can always add a constant to the Hamiltonian without changing the physics. But that's not the case for supersymmetric quantum mechanics (nor, indeed, for supersymmetric quantum field theories). The requirement that  $E \geq 0$  also rules out some very familiar quantum mechanical potentials, like  $V = -1/r$  of the hydrogen atom. The potential in supersymmetric quantum mechanics must always be positive definite.

As an aside: there's only one other place in physics where we care about the overall value of the ground state energy, and that's the cosmological constant in general relativity. So far, sadly, no plausible link has been found between the value of the cosmological constant and the supersymmetry algebra.

We can learn more from the supersymmetry algebra. The energy eigenstates of supersymmetric quantum mechanics are *almost* always degenerate. Consider the set of states with some fixed energy  $E$ ,

$$H|\psi\rangle = E|\psi\rangle$$

It's simple to check from the supersymmetry algebra (1.1) that  $[H, Q] = [H, Q^\dagger] = 0$ , facts which require us to also use  $Q^2 = Q^{\dagger 2} = 0$ . This means that the operators  $Q$  and  $Q^\dagger$  act within an energy eigenspace. If the energy is  $E \neq 0$ , we have

$$\{Q, Q^\dagger\} = 2E \quad \Rightarrow \quad \{c, c^\dagger\} = 1 \quad \text{with } c = \frac{Q}{\sqrt{2E}} \quad (1.3)$$

We also have  $c^2 = c^{\dagger 2} = 0$ . This is the same algebra that is formed by fermionic creation and annihilation operators. The algebra has a two-dimensional irreducible representation spanned by the states  $|0\rangle$  and  $|1\rangle$  with the properties that

$$c|0\rangle = 0 \quad \text{and} \quad |1\rangle = c^\dagger|0\rangle$$

Equivalently we have  $c^\dagger|1\rangle = 0$  and  $|0\rangle = c|1\rangle$ . The algebra is telling us that all energy eigenstates with  $E \neq 0$  states must come in pairs. Of course, there could be a still bigger degeneracy, with several pairs all having the same energy. But, at each energy level, the number of states must be even.

The one exception is when we have states with energy  $E = 0$ . As we've seen, if such states exist then they are necessarily the ground states. Importantly, the argument above that enforces the degeneracy of the spectrum fails: it is quite possible to have a lone ground state  $|\Omega\rangle$  because, as we can see in (1.2), any such ground state necessarily obeys  $Q|\Omega\rangle = Q^\dagger|\Omega\rangle = 0$ . Again, it's quite possible to have more than one ground state. But if that's the case, they're not related by the action of  $Q$  or  $Q^\dagger$ .

Finally, there is a slightly more formal way of viewing the story above. Inspired by the connection to fermionic creation operators, we define the "fermion number operator"

$$F = c^\dagger c$$

This obeys  $[F, Q] = -Q$  and  $[F, Q^\dagger] = Q^\dagger$  and  $[F, H] = 0$ . Clearly this operator is well defined only on states with energy  $E \neq 0$ , where it acts as  $F|0\rangle = 0$  and  $F|1\rangle = |1\rangle$ .

Correspondingly, the Hilbert space decomposes into “bosonic states” with  $F = 0$  and “fermionic states” with  $F = 1$ ,

$$\mathcal{H} = \mathcal{H}_B \oplus \mathcal{H}_F \tag{1.4}$$

We say that there is a  $\mathbb{Z}_2$  *grading* of the Hilbert space. The  $E \neq 0$  pairs have one state in  $\mathcal{H}_B$  and one in  $\mathcal{H}_F$ . As it stands, it’s not clear which of these Hilbert spaces we should assign the  $E = 0$  states to. This will become clearer when we turn to specific examples below.

Finally, one last piece of terminology. If a ground state with energy  $E = 0$  exists, then we say that supersymmetry is *unbroken*. If the ground state has energy  $E > 0$  then we say that supersymmetry is *broken*. This language is really adopted from higher dimensions where symmetries that do not leave the vacuum invariant are said to be “spontaneously broken”. In the present context we say that supersymmetry is broken if the vacuum is not annihilated by the supercharges: the connection to symmetries will become clearer as we proceed.

## 1.2 A Particle in a Potential

An abstract algebra like (1.1) is all well and good, but to build intuition we really need a concrete example that realises this algebra. Happily such an example exists: we consider the quantum mechanics of a particle moving on a line. The only small novelty is that the particle has an internal degree of freedom, like spin, that can take two different values. The Hilbert space is

$$\mathcal{H} = L^2(\mathbb{R}) \otimes \mathbb{C}^2$$

where the  $L^2(\mathbb{R})$  means normalisable functions on the real line  $\mathbb{R}$  which, of course, is simply the Hilbert space for a particle on a line. Meanwhile the  $\mathbb{C}^2$  factor is the internal degree of freedom. In keeping with the notation of the previous section, we’ll take the internal states of the  $\mathbb{C}^2$  factor to be spanned by  $|0\rangle$  and  $|1\rangle$ . The Hilbert space then decomposes into our “fermionic” and “bosonic” pieces,

$$\mathcal{H} = L^2(\mathbb{R})|0\rangle \oplus L^2(\mathbb{R})|1\rangle = \mathcal{H}_B \oplus \mathcal{H}_F$$

In this context, it might be better to think of  $|0\rangle$  and  $|1\rangle$  as a spin degree of freedom, with  $\mathcal{H}_B$  and  $\mathcal{H}_F$  the “spin down” and “spin up” components of the Hilbert space. On the other hand, it might be confusing to think of “spin up” as a fermion and “spin down” as a boson so I stress that these are just names at this stage and don’t come with any other fermionic/bosonic connotations. We’ll use both pieces of terminology in what follows.

For our supercharge  $Q$ , we take

$$Q = (p - ih'(x)) \otimes \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \quad (1.5)$$

Here  $p = -id/dx$  is the usual momentum operator (in units where  $\hbar = 1$ ) and  $h(x)$  is a real function. We have  $Q^2 = 0$  because the  $2 \times 2$  matrix squares to zero. Taking the conjugate gives

$$Q^\dagger = (p + ih'(x)) \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \quad (1.6)$$

and so

$$H = \frac{1}{2}(QQ^\dagger + Q^\dagger Q) = \frac{1}{2}(p^2 + h'^2) \mathbb{1} - \frac{1}{2}h''\sigma^3 \quad (1.7)$$

The first factor is the familiar Hamiltonian for a particle with unit mass moving on a line with potential

$$V(x) = \frac{1}{2} \left( \frac{dh}{dx} \right)^2 \quad (1.8)$$

This term comes with the  $2 \times 2$  unit matrix  $\mathbb{1}$  and so doesn't care about the spin of the particle. In contrast, the second term comes with the Pauli matrix

$$\sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

This term acts like a magnetic field, distinguishing spin up and spin down by the minus sign.

The operator  $F$  that distinguishes spin up from spin down is simply

$$F = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad (1.9)$$

This tells us that the ‘‘bosonic’’ or ‘‘spin down’’ part of the Hilbert space  $\mathcal{H}_B$  is composed of states of the form  $\psi(x)|0\rangle = \psi(x)\begin{pmatrix} 0 \\ 1 \end{pmatrix}$  and the ‘‘fermionic’’ or ‘‘spin up’’ part of the Hilbert space  $\mathcal{H}_F$  is composed of states of the form  $\psi(x)|1\rangle = \psi(x)\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ . Note that the definition of  $F$  now happily extends to the zero energy states as well.

### 1.2.1 Ground States

Usually, it is challenging to find the exact ground states of any quantum mechanical potential. One of the rather pretty features of supersymmetric quantum mechanics is that we can sometimes find exact expressions for the ground states.

To kick things off, let's look at the semi-classical ground states. The potential energy (1.8) is positive definite and has a minimum whenever there is a critical point of  $h$ ,

$$V(x) = 0 \quad \Leftrightarrow \quad h'(x) = 0$$

If we Taylor expand around such a critical point  $x = x_0$ , we have

$$h(x) \approx h(x_0) + \frac{1}{2}\omega(x - x_0)^2 + \dots$$

This gives a potential energy (1.8) that is, to leading order, a harmonic oscillator,  $V(x) = \frac{1}{2}\omega^2(x - x_0)^2 + \dots$ . While the classical ground state energy of a harmonic oscillator vanishes, quantum mechanically we have  $E_0 = \frac{1}{2}|\omega|$  (working in units with  $\hbar = 1$ .) But the supersymmetric system also gets a contribution only from the spin-dependent term in (1.7) which, at leading order, is

$$\Delta E = \pm \frac{1}{2}|\omega| \tag{1.10}$$

If we take the minus sign, this precisely cancels the contribution from the harmonic oscillator ground state energy, giving us a total, semi-classical energy  $E = 0$ . This simple minded analysis shows that it's quite plausible that zero energy ground states exist in this system.

Let's now look more closely at the full quantum problem and, in particular, the question of whether  $E = 0$  ground states exist. A general state takes the form

$$\Psi(x) = \begin{pmatrix} \psi(x) \\ \phi(x) \end{pmatrix} \tag{1.11}$$

But to qualify as an  $E = 0$  ground state, this must be annihilated by both the supercharges,  $Q\Psi = Q^\dagger\Psi = 0$ , meaning that

$$-i \left( \frac{d}{dx} + h' \right) \psi = 0 \quad \text{and} \quad -i \left( \frac{d}{dx} - h' \right) \phi = 0$$

The magic of supersymmetry means that, at least for the ground state, the Schrödinger equation has morphed from a challenging second order differential equation into a pair of decoupled, first order differential equations. Note that this same trick doesn't work to figure out the excited states of the theory. We can't solve for the whole spectrum. But we can solve for the ground state.

Indeed, the equations are straightforward to solve. We have

$$\psi(x) = e^{-h} \quad \text{and} \quad \phi(x) = e^{+h} \quad (1.12)$$

There is, as always in quantum mechanics, one last criterion: we need to determine if these states are normalisable. This clearly depends on the form of  $h(x)$  which, in turn, determines the potential energy (1.8). There are three possibilities

- If  $h \rightarrow +\infty$  as  $|x| \rightarrow +\infty$  then  $\psi(x)$  is normalisable and we must have  $\phi(x) = 0$ . In this case there is a unique ground state that sits in the “fermionic” or “spin up” part of the Hilbert space  $\mathcal{H}_F$ .
- If  $h \rightarrow -\infty$  as  $|x| \rightarrow +\infty$  then  $\phi(x)$  is normalisable and we must have  $\psi(x) = 0$ . In this case there is a unique ground state that sits in the “bosonic” or “spin down” part of the Hilbert space  $\mathcal{H}_B$ .
- If  $h$  has neither of these properties, then there is no  $E = 0$  ground state and supersymmetry is broken. In this case the ground state necessarily has  $E \neq 0$  and is degenerate.

To get a better sense of what’s going on, let’s look at some simple examples.

#### Example 1: Quadratic $h$


To start, we take  $h = \frac{1}{2}\omega x^2$  with  $\omega > 0$ . In this case, we just have a harmonic oscillator with potential energy (1.8) given by  $V(x) = \frac{1}{2}\omega^2 x^2$ . The additional spin-dependent term in the Hamiltonian (1.7) just shifts the spectrum up or down by  $\frac{1}{2}\omega$ . The upshot is that the “fermionic” or “spin up” spectrum in  $\mathcal{H}_F$  takes the form

$$E_F = \omega n \quad n = 0, 1, 2, \dots$$

Here we find the unique ground state. Meanwhile, the “bosonic” or “spin down” spectrum in  $\mathcal{H}_B$  takes the form

$$E_B = \omega n \quad n = 1, 2, \dots$$

As promised, all excited states with  $E > 0$  are paired, but there is a single unpaired ground state at  $E = 0$ .



**Figure 1.** The potential for a cubic  $h$  has two classical  $E = 0$  ground states.

Note that, had we chosen  $\omega < 0$ , the situation would be reversed with the ground state living in  $\mathcal{H}_B$ .


### Example 2: Cubic $h$

When  $h$  is a polynomial of degree higher than two, we can't solve the entire spectrum. But we can get a good understanding of the ground states. Suppose that we take

$$h = \lambda x^3 + \dots$$


where  $\dots$  are lower order monomials. As we have seen, in this case there can be no  $E = 0$  ground states.

A typical form of  $h$  is shown on the left of Figure 1, with the corresponding potential  $V(x)$  on the right. If we neglected the spin degree of freedom, we would have the familiar double well potential of quantum mechanics and we have some intuition about what happens in this case. Clearly there are two classical minima with  $V(x) = 0$  and we can construct



an approximation to the ground state with a Gaussian wavefunction that is localised at one, or other, of the minima, as shown on the right, with the orange and green curves each showing different candidate ground state wavefunctions.

Since  $h'' < 0$  near the left-hand minimum we expect that this wavefunction can lower its energy by sitting in the “spin down” part of the Hilbert space  $\mathcal{H}_B$ . Similarly,  $h'' > 0$



**Figure 2.** The potential for a quartic  $h$  has three classical  $E = 0$  ground states.

near the right-hand minimum so we expect that it's energetically preferable for this wavefunction to sit in  $\mathcal{H}_F$ .

Usually in a double well potential, the particle can lower its energy by tunnelling through the barrier and sitting in a superposition of both states. But that's not the case here because the two wavefunctions live in different components of spin space. This kills the possibility for tunnelling. Instead, the supersymmetric set-up is closer to our naive, classical guess of the ground states, with a Gaussian around each minima giving a good approximation to the ground state. Our arguments above tell us that the energy of this two-fold degenerate ground state is necessarily  $E > 0$ . We will say more about tunnelling in this system and how to compute the actual energy in Section 2.2.


### Example 3: Quartic $h$

Next consider  $h(x)$  of the form

$$h = \lambda x^4 + \dots \tag{1.13}$$

where  $\dots$  are terms of order cubic and lower. We pick  $\lambda > 0$  so that  $h \rightarrow +\infty$  as  $|x| \rightarrow \infty$ .

A typical  $h(x)$  and the associated potential  $V(x)$  are shown in Figure 2. There are now three classical ground states and a naive semi-classical approach would suggest that we can approximate the true ground states as Gaussians localised around any of these three minima. The two outside minima have  $h'' > 0$  and so the lowest energy wavefunctions live in  $\mathcal{H}_F$ , while the middle minimum has  $h'' < 0$  and so the lowest energy state sits in  $\mathcal{H}_B$ .



**Figure 3.** The exact  $E = 0$  wavefunction is localised only on the outer minima.

However, this time we know the exact ground state: it is given by  $\psi(x) = e^{-h(x)}$  and lives in  $\mathcal{H}_F$ . This is plotted in orange, superposed on the potential, in Figure 3. The wavefunction is peaked on those places where  $h < 0$  which, in this case, means that two outer minima. This clearly demonstrates the tunnelling phenomena, in which the true ground state sits in a superposition of minima but, as you can see, there is not necessarily a symmetric distribution between the two vacua.

We started with three states that we thought had the smallest energy – one for each minima – but only one survives as the true  $E = 0$  ground state. The other two states must have some small, but non-zero energy. These states are the Gaussian localised in the middle vacuum, and the combination of states localised on the outside minima that is orthogonal to the ground state. Although it is far from obvious from staring at the potential, supersymmetry tells us that the energies of these states must be degenerate.

As we vary the parameters in the function  $h(x)$ , the energy spectrum of the theory will change. However, the energy of the ground state remains pinned to  $E = 0$ . The one exception to this statement occurs if we sent  $\lambda \rightarrow 0$ . In this case, one of the minima of the potential runs off to infinity, as  $x \sim 1/\lambda$ , and carries the  $E = 0$  ground state wavefunction with it. In this case, we go over to the situation of a cubic  $h(x)$  described above in which there are two ground states, both with  $E > 0$ .

### 1.2.2 The Witten Index

The robustness of supersymmetric ground states can be formulated more generally using the *Witten index*. As we've seen, the Hilbert space decomposes into two pieces

$$\mathcal{H} = \mathcal{H}_B \oplus \mathcal{H}_F$$

These two pieces are characterised by the “fermion number operator”  $F$  which has eigenvalues 0 or 1. It is often more useful to consider the operator  $(-1)^F$ , sometimes called *fermion parity*, that takes eigenvalues  $+1$  on states in  $\mathcal{H}_B$  and  $-1$  on states in  $\mathcal{H}_F$ . For our example of a particle on a line, it's simple to check that  $(-1)^F = -\sigma^3$ .

The Witten index is defined as

$$\mathcal{I} = \text{Tr} (-1)^F e^{-\beta H}$$

Here the trace is taken over all states in the Hilbert space. The parameter  $\beta$  plays a role like inverse temperature  $\beta = 1/k_B T$  in statistical mechanics. The Witten index differs from the usual statistical mechanical partition function by the signs  $(-1)^F$ . Importantly, as we will now argue, in supersymmetric theories the Witten index is actually independent of  $\beta$

$$\frac{d\mathcal{I}}{d\beta} = 0$$

This follows because, as we have seen, the spectrum of supersymmetric quantum mechanics is degenerate for any state with  $E > 0$ . Formally, there is an isomorphism between  $\mathcal{H}_B$  and  $\mathcal{H}_F$ ,

$$\mathcal{H}_B \Big|_{E>0} \cong \mathcal{H}_F \Big|_{E>0}$$

This means that the trace over any state with  $E > 0$  simply cancels out in the Witten index: for every  $+e^{-\beta E}$  from  $\mathcal{H}_B$  there is a corresponding  $-e^{-\beta E}$  from  $\mathcal{H}_F$ . This means that the Witten index only receives contributions from the zero energy states which, as we've seen, need not be duplicated in both  $\mathcal{H}_B$  and  $\mathcal{H}_F$ . In other words, the Witten index really counts the difference in the number of ground states in each sector,

$$\mathcal{I} = \dim \mathcal{H}_{0,B} - \dim \mathcal{H}_{0,F}$$

where  $\mathcal{H}_{0,B}$  is the space of  $E = 0$  bosonic ground states, and similar for  $\mathcal{H}_{0,F}$ .

Before we proceed, a few comments. Since  $\mathcal{I}$  doesn't depend on  $\beta$ , you might wonder why we don't just set  $\beta = 0$  and consider  $\text{Tr}(-1)^F$ . Indeed, often the Witten index is written in this way as shorthand, but it's a dangerous thing to do. The quantity  $\text{Tr}(-1)^F$  is an infinite series of  $+1$  and  $-1$  and by pairing terms together in various ways you can get any answer that you like. Including  $e^{-\beta H}$  in the definition acts as a regulator for this sum, rendering it finite. Of course, it's a familiar regulator because it also appears in the partition function in statistical mechanics.


The same arguments that show  $d\mathcal{I}/d\beta = 0$  also show that  $\mathcal{I}$  is independent of the parameters of the Hamiltonian  $H$ . This was demonstrated in the examples above although, as we also saw, it comes with a caveat: if you change the Hamiltonian too dramatically then you can lose states in your Hilbert space and this will change  $\mathcal{I}$ . This happens for the particle on a line whenever we change the power of the leading term in  $h(x)$ .

The Witten index counts the difference between the bosonic and fermionic  $E = 0$  states. However, in the simple examples considered above, it actually counts the number of  $E = 0$  states, positive if they're bosonic, negative if they're fermionic. One might wonder if, in practice, it always does this. Indeed, there is some intuition that suggests this is the case. If there's no good reason for pairs of states to be stuck at  $E = 0$  then, as you vary parameters in the potential, it's tempting to think that they will be lifted to  $E > 0$ .

However, it's not difficult to exhibit examples where, for example,  $\mathcal{I} = 0$  but there are a pair of bosonic and fermionic  $E = 0$  states. A particularly simple example arises from particle moving on a circle  $\mathbf{S}^1$  of radius  $R$ . The supercharge (1.5) and Hamiltonian (1.7) take the same form as before and are characterised by a periodic function  $h(x) = h(x + 2\pi R)$ . We can follow our earlier footsteps to find a two parameter family of ground states labelled by  $\alpha, \beta \in \mathbb{C}$ ,

$$\Psi(x) = \alpha \begin{pmatrix} e^{-h} \\ 0 \end{pmatrix} + \beta \begin{pmatrix} 0 \\ e^{+h} \end{pmatrix}$$

This time, because the particle lives on a circle, there is no issue with the normalisability of the wavefunction. We see that the system has two linearly independent  $E = 0$  ground states for any choice of  $h$ . Yet, because one ground state lives in  $\mathcal{H}_B$  and the other in  $\mathcal{H}_F$ , the Witten index of this system is  $\mathcal{I} = 0$ . The potential (in blue) and wavefunctions (in orange and green) for  $h(x) = \sin(x/R)$  are shown in Figure 4.



**Figure 4.** Two  $E = 0$  ground states for the double well potential on a circle.

For this particle on the circle, the pair of states sticks at  $E = 0$  as we change the parameters of  $h$ , even though these ground states are not protected by the Witten index. One might wonder if there's a deeper reason for this. There is and it's related to the deeper mathematical concept of *cohomology*. We'll look at this further in Section 3.

Finally, one last comment before we move on. The manipulations of the Witten index rely on the discreteness of the energy spectrum. There are more subtle situations, where a particle moves on a non-compact space without a potential, where the energy spectrum is continuous and, despite the bose-fermi degeneracy in the spectrum, strange things can happen that mean that  $\mathcal{I}$  does, in fact, depend on  $\beta$ . We will not encounter situations of this kind in these lectures.

### 1.3 The Supersymmetric Action

There is one fairly large omission in our discussion so far. As presented above, supersymmetric Hamiltonians have a nice algebraic structure. But we have no inkling of why supersymmetry has anything to do with symmetry!

A clue is to be found in the commutation relations

$$[H, Q] = [H, Q^\dagger] = 0 \quad \Rightarrow \quad [H, Q + Q^\dagger] = 0$$

Usually in quantum mechanics, Hermitian operators that commute with the Hamiltonian correspond to conserved quantities and conserved quantities come, via Noether's theorem, from symmetries. This suggests that perhaps  $Q + Q^\dagger$  is somehow the conserved charge associated to a symmetry. But what symmetry?

Often the Lagrangian framework is a better starting point when looking for symmetries. To this end, we would like to introduce a Lagrangian for our supersymmetric theory of a particle on a line. We know well how to think of position and momentum in the Lagrangian setting. But how do we incorporate the discrete  $\mathbb{C}^2$  factor in the Hilbert space that gave us the all-important  $\mathbb{Z}_2$  grading?

The answer is that we should turn to fermions. In higher dimensions, adding a fermion to a Lagrangian gives another field. But in quantum mechanics, fermions simply offer a different way of describing some discrete aspect of the physics.

To illustrate this, consider the action

$$S = \int dt L = \int dt \left[ \frac{1}{2} \dot{x}^2 + i\psi^\dagger \dot{\psi} - \frac{1}{2} h'^2 + h'' \psi^\dagger \psi \right] \quad (1.14)$$

where  $\psi$  and its conjugate  $\psi^\dagger$  are Grassmann variables. Note that their kinetic terms are first order, like the Dirac action that we met in [Quantum Field Theory](#), albeit without the intricacies of gamma matrices. We will first show that this action is equivalent to the supersymmetric Hamiltonian (1.7) describing a particle with an internal degree of freedom moving on a line. We'll then understand how to think of the supercharges  $Q$  in the Lagrangian formulation.

To construct the Hamiltonian from a Lagrangian, we proceed in the usual manner. We first introduce the conjugate momentum for both bosonic and fermionic degrees of freedom

$$p(t) = \frac{\delta S}{\delta \dot{x}(t)} = \dot{x}(t) \quad \text{and} \quad \pi(t) = \frac{\delta S}{\delta \dot{\psi}(t)} = -i\psi^\dagger(t)$$

In the quantum theory, these obey the canonical (anti)-commutation relations

$$[x, p] = i \quad \text{and} \quad \{\psi, \psi^\dagger\} = 1 \quad (1.15)$$

which, in the Heisenberg picture, hold at a fixed time  $t$ . The Hamiltonian is then the Legendre transform

$$H = \dot{x}p + \dot{\psi}\pi - L$$

There is, however, a small subtlety awaiting us. We think of the Lagrangian as a classical object in which  $x$  and  $\dot{x} = p$  be placed in any order. Relatedly,  $\psi$  and  $\psi^\dagger$  are viewed as “classical Grassmann variables” in the action, which means that if one moves past the other then we just pick up a minus sign. But in the Hamiltonian, these are all to be thought of as quantum operators and, because of the commutation relations (1.15), ordering matters. Which ordering should we take?

This kind of ordering ambiguity is not uncommon when going from classical to quantum systems. In the present situation we don't have to worry about  $x$  and  $p$  (although we will later in these lectures) but only about the ordering of  $\psi$  and  $\psi^\dagger$ . In the action, it doesn't matter whether we write the last term as  $h''\psi^\dagger\psi$  or  $-h''\psi\psi^\dagger$ : they are the same. But in the Hamiltonian, they differ by a constant because, when viewed as quantum operators,  $\psi^\dagger\psi = -\psi\psi^\dagger + 1$ . In most other contexts, there is no way to fix this ambiguity and it reflects the fact that there are different ways to quantise a classical theory. However, for us, we do have a way to fix the ambiguity since the resulting Hamiltonian should be supersymmetric. The correct answer, as we will see, is to take

$$H = \frac{1}{2}(p^2 + h'^2) - \frac{1}{2}h''(\psi^\dagger\psi - \psi\psi^\dagger) \quad (1.16)$$

where, in the final term, we've split the difference and treated  $\psi^\dagger\psi$  and  $\psi\psi^\dagger$  in a symmetric fashion.

To make contact with our previous notation, we just need to appreciate that, due to their Grassmann nature,  $\{\psi, \psi\} = \{\psi^\dagger, \psi^\dagger\} = 0$  which, in conjunction with the anti-commutation relation (1.15), has a two-dimensional real representation. Indeed, we met this before when discussing the energy spectrum of supersymmetric quantum mechanics in (1.3). The representation can be thought of as simply replacing the Grassmann variables with  $2 \times 2$  matrices,

$$\psi \rightarrow \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad \psi^\dagger \rightarrow \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

This then gives  $\psi^\dagger\psi - \psi\psi^\dagger = \sigma^3$ , and the Hamiltonian (1.16) coincides with our previous result (1.7).

Written in terms of the fermionic degrees of freedom, the supercharges (1.5) and (1.6) take the form

$$Q = (p - ih')\psi \quad \text{and} \quad Q^\dagger = (p + ih')\psi^\dagger \quad (1.17)$$

The (anti)-commutation relations of  $Q$  with the various fields are

$$\begin{aligned} [Q, x] &= -i\psi & , & & [Q^\dagger, x] &= -i\psi^\dagger \\ \{Q, \psi\} &= 0 & , & & \{Q^\dagger, \psi\} &= p + ih' \\ \{Q, \psi^\dagger\} &= p - ih' & , & & \{Q^\dagger, \psi^\dagger\} &= 0 \end{aligned} \quad (1.18)$$

You can check that these commutation relations give  $\{Q, Q^\dagger\} = 2H$ , with  $H$  given in (1.16), as they should.

### 1.3.1 Supersymmetry as a Fermionic Symmetry

Now we can see what this has to do with symmetry. The action (1.14) has the special property that it is invariant under the following *supersymmetry transformations*

$$\begin{aligned}\delta x &= \epsilon^\dagger \psi - \epsilon \psi^\dagger \\ \delta \psi &= \epsilon(-i\dot{x} + h') \\ \delta \psi^\dagger &= \epsilon^\dagger(i\dot{x} + h')\end{aligned}\tag{1.19}$$

Note that these swap bosonic fields  $x$  for fermionic fields  $\psi$ . This is the characteristic feature of supersymmetry that distinguishes it from other symmetries. For this to make sense, the infinitesimal transformation parameter  $\epsilon$  must be a Grassmann valued object.

Let's first check that the action (1.14) is indeed invariant under the supersymmetry transformations as claimed. A generic variation of the action gives

$$\delta S = \int dt \left[ \dot{x} \delta \dot{x} + i \delta \psi^\dagger \dot{\psi} + i \psi^\dagger \delta \dot{\psi} - h' h'' \delta x + h''' \delta x \psi^\dagger \psi + h'' (\delta \psi^\dagger \psi + \psi^\dagger \delta \psi) \right]$$

Now we substitute in the particular supersymmetry transformation (1.19). We collate the  $\epsilon$  and  $\epsilon^\dagger$  terms on different lines to find

$$\begin{aligned}\delta S &= \int dt \epsilon \left[ -\dot{x} \psi^\dagger - i \psi^\dagger \frac{d}{dt} (-i\dot{x} + h') + h' h'' \psi^\dagger - h''' \psi^\dagger \psi^\dagger \psi - h'' \psi^\dagger (-i\dot{x} + h') \right] \\ &\quad + \epsilon^\dagger \left[ \dot{x} \psi + i(i\dot{x} + h') \dot{\psi} - h' h'' \psi + h''' \psi \psi^\dagger \psi + h'' (i\dot{x} + h') \psi \right]\end{aligned}$$

There are some minus signs to ensare the unwary: these arise in moving the  $\epsilon$  parameters past other Grassmann objects.

We can immediately discard many terms. First, the  $\dot{x} \psi$  and  $\dot{x} \psi^\dagger$  terms cancel (for the latter, after an integration by parts). Second the  $h'''$  terms disappear on Grassmann grounds. We're left with

$$\begin{aligned}\delta S &= \int dt \epsilon \left[ i \psi^\dagger h' + h' h'' \psi^\dagger - h'' \psi^\dagger (-i\dot{x} + h') \right] \\ &\quad + \epsilon^\dagger \left[ i h' \dot{\psi} - h' h'' \psi + h'' (i\dot{x} + h') \psi \right] \\ &= \int dt i \epsilon \frac{d}{dt} (h' \psi^\dagger) + i \epsilon^\dagger \frac{d}{dt} (h' \psi)\end{aligned}$$

But this is a total derivative and so we have

$$\delta S = 0\tag{1.20}$$

as advertised.

Before we go on, it will be useful to present this result in a slightly different way. We can think of the transformations (1.19) as generated by the following fermionic operators

$$\begin{aligned} \mathcal{Q} &= \int dt \left[ \psi(t) \frac{\delta}{\delta x(t)} + (i\dot{x} + h') \frac{\delta}{\delta \psi^\dagger(t)} \right] \\ \mathcal{Q}^\dagger &= \int dt \left[ -\psi^\dagger(t) \frac{\delta}{\delta x(t)} - (i\dot{x} - h') \frac{\delta}{\delta \psi(t)} \right] \end{aligned} \quad (1.21)$$

Here, the functional derivatives act as

$$\frac{\delta}{\delta x(t)} x(t') = \delta(t - t') \quad \text{and} \quad \frac{\delta}{\delta x(t)} \psi(t') = \frac{\delta}{\delta x(t)} \psi^\dagger(t') = 0$$

with similar expressions for the fermions. The supersymmetry transformations (1.19) can then be written as, for example,  $\delta x(t) = (\epsilon^\dagger \mathcal{Q} + \epsilon \mathcal{Q}^\dagger)x(t)$ . The invariance of the action (1.20) becomes simply

$$\mathcal{Q}S = \mathcal{Q}^\dagger S = 0 \quad (1.22)$$

This form of the generators will be useful in Section 2 when we discuss the path integral formulation of supersymmetric quantum mechanics.

### The Supercharge is a Noether Charge

Finally, we can make good on our promise and see that the supercharges  $Q$  and  $Q^\dagger$  are indeed Noether charges for supersymmetry. Usually when the action has a symmetry, we can construct the Noether charge by allowing the transformation parameter to depend on time. Things are no different here. We vary the action with  $\epsilon = \epsilon(t)$ . There are two steps where things differ from our previous calculation: first when we vary the kinetic terms, and again at the last where we see that the variation of the Lagrangian is a total derivative which requires an integration by parts. We end up with

$$\delta S = \int dt \dot{\epsilon}^\dagger Q$$

where the Noether charge  $Q$  in this calculation coincides with our previous expression (1.17) for the supercharge:  $Q = (\dot{x} - ih')\psi = (p - ih')\psi$ .

It's slightly odd that the variation of the action involves  $\dot{\epsilon}^\dagger$  but not  $\dot{\epsilon}$ . We can trace this to our choice of fermion kinetic term  $\psi^\dagger \dot{\psi}$ , which is asymmetric between  $\psi$  and  $\psi^\dagger$ . We could instead start with the more symmetric choice

$$S = \int dt L = \int dt \left[ \frac{1}{2} \dot{x}^2 + \frac{i}{2} (\psi^\dagger \dot{\psi} - \dot{\psi}^\dagger \psi) - \frac{1}{2} h'^2 + h'' \psi^\dagger \psi \right]$$

Clearly this is equivalent to our original action (1.14) after an integration by parts, but it's sometimes best to work with these kinds of symmetric kinetic terms for fermions, and computing the Noether charge is one such place. We now find

$$\delta S = \int dt \frac{1}{2} \epsilon^\dagger \dot{Q} - \frac{1}{2} \dot{\epsilon} Q^\dagger$$

where  $Q^\dagger = (p + ih')\psi^\dagger$ .

We can now go full circle. In the operator framework of quantum mechanics, the Noether charges generate the symmetry. Again, supersymmetry is no different. The transformation of any field is given as

$$\delta(\text{field}) = i[\epsilon^\dagger Q - \epsilon Q^\dagger, \text{field}]$$

where, as before,  $\epsilon$  is a Grassmann valued parameter. The minus sign in the expression above ensures that  $\epsilon^\dagger Q - \epsilon Q^\dagger$  is Hermitian (because  $(\epsilon^\dagger Q)^\dagger = Q^\dagger \epsilon = -\epsilon Q^\dagger$ ) and the overall factor of  $i$  ensures that  $\delta x$  is Hermitian. Using the commutation relations (1.18), you can check that we recover the supersymmetry transformations (1.19) as promised.

## 1.4 A Particle Moving in Higher Dimensions

There are some straightforward generalisations of the supersymmetric theories of a particle moving on a line that we considered in the last section. These will bring out a number of new themes that we will return to as these lectures progress.

### 1.4.1 A First Look at Morse Theory

We start with a direct generalisation of our earlier supersymmetric system to a particle moving in  $\mathbb{R}^n$ , parameterised by coordinates  $x^i$  with  $i = 1, \dots, n$ .

The observation that supersymmetry relates bosonic to fermionic fields suggests that we should also introduce  $n$  Grassmann valued fields  $\psi^i$ , with  $i = 1, \dots, n$ . These obey the anti-commutation relations

$$\{\psi^i, \psi^j\} = \{\psi^{\dagger i}, \psi^{\dagger j}\} = 0 \quad \text{and} \quad \{\psi^i, \psi^{\dagger j}\} = \delta^{ij} \quad (1.23)$$

As in our previous discussion, these fermionic fields should be viewed as operators acting on some internal, finite dimensional Hilbert space. To construct a representation we introduce the ‘‘Fock vacuum’’ state  $|0\rangle$  that obeys

$$\psi^i |0\rangle = 0 \quad \text{for all } i = 1, \dots, n$$

We can then build up the space of states by acting on  $|0\rangle$  with  $\psi^{\dagger i}$ , recalling that you only get to act with a given  $\psi^{\dagger}$  once. This means that the spectrum of internal states takes the form

$$\begin{aligned} &|0\rangle \\ &\psi^{\dagger i}|0\rangle \\ &\psi^{\dagger i}\psi^{\dagger j}|0\rangle \\ &\vdots \\ &\psi^{\dagger 1}\dots\psi^{\dagger n}|0\rangle \end{aligned}$$

There are  $\binom{n}{p}$  states in the sector where we act with  $p$  different  $\psi^{\dagger}$ 's. The total number of states is then

$$\sum_{p=0}^n \binom{n}{p} = (1+1)^n = 2^n$$

where you should expand out the  $(1+1)^n$  in the middle using the binomial theorem to get the sum on the left. This means that our supersymmetric quantum mechanics will describe a particle moving in  $\mathbb{R}^n$  with  $2^n$  internal states.

There's a useful geometrical way to think about these states. At the top of the pyramid depicted above we have wavefunctions that look like  $\phi(x)|0\rangle$ : these are just functions over  $\mathbb{R}^n$ .

At the next level, the wavefunctions look like  $\phi(x)\psi^{\dagger i}|0\rangle$  and come with an internal index  $i = 1, \dots, n$ . We usually think of objects on  $\mathbb{R}^n$  that carry such an index as vectors. However, as we now explain, the anti-symmetric nature of the Grassmann variable means that it's much more natural to think about these states as one-forms on  $\mathbb{R}^n$ .

We really see why it's useful to think of these states as forms when we get to the second level. Here wavefunctions look like  $\phi(x)\psi^{\dagger i}\psi^{\dagger j}|0\rangle = -\phi(x)\psi^{\dagger j}\psi^{\dagger i}|0\rangle$ , with the  $i, j$  index necessarily anti-symmetric. But this is precisely the definition of a two-form in differential geometry. This then continues until we reach the unique top form  $\psi^{\dagger 1}\dots\psi^{\dagger n}|0\rangle$ . All of this suggests that we should make the identification between Grassmann variables and forms

$$\psi^{\dagger i} \longleftrightarrow dx^i \wedge$$

On  $\mathbb{R}^n$ , there's little advantage to be had in working with  $p$ -forms rather than just sticking with Grassmann variables and, for the rest of this section, we'll use the latter

notation. However, this relationship to  $p$ -forms will be of crucial importance when we turn to more geometrical settings in Section 3.

The supersymmetric quantum mechanics also has a fermion parity operator  $(-1)^F$  which simply counts the number of excited fermions mod 2. By convention, we take  $F|0\rangle = 0$  so  $(-1)^F|0\rangle = +1$ . Then if  $|p\rangle$  denotes a state in the sector with  $p$  excited fermions, we have

$$(-1)^F|p\rangle = (-1)^p|p\rangle \quad (1.24)$$

In other words,  $(-1)^F$  counts the degree of a  $p$ -form, mod 2.

### The Supersymmetric Hamiltonian

The supersymmetric quantum mechanics for a particle moving in  $\mathbb{R}^N$  involves a real function  $h(x^i)$  and the Hamilton

$$H = \frac{1}{2} \sum_{i=1}^n (p_i^2 + (\partial_i h)^2) - \frac{1}{2} \partial_i \partial_j h [\psi^{\dagger i}, \psi^j] \quad (1.25)$$

It's not difficult to check that this can be written in the defining way  $H = \frac{1}{2}\{Q, Q^\dagger\}$  with the standard (anti)-commutation relations and the supercharges

$$Q = (p_i - i\partial_i h)\psi^i \quad \text{and} \quad Q^\dagger = (p_i + i\partial_i h)\psi^{\dagger i} \quad (1.26)$$

where summation convention is used, both for the supercharges and the final term of the Hamiltonian.

We can compute the Witten index by looking at the semi-classical ground states. The bosonic part of the Hamiltonian has a ground state at any critical point  $x = X$ ,

$$\partial_i h(X) = 0 \quad \text{for all } i = 1, \dots, n$$

But where does this ground state sit in the internal space? First recall what happened in the simpler case where we just had one fermion and, correspondingly, two states  $|0\rangle$  and  $|1\rangle$  with  $\psi|0\rangle = 0$  and  $|1\rangle = \psi^\dagger|0\rangle$ . In that case, the final term in the Hamiltonian was

$$H_{\text{Fermi}} = -\frac{1}{2}h''[\psi^\dagger, \psi]$$

So acting on the two states, we had

$$\begin{aligned} H_{\text{Fermi}}|0\rangle &= +\frac{1}{2}h''|0\rangle \\ H_{\text{Fermi}}|1\rangle &= -\frac{1}{2}h''|1\rangle \end{aligned}$$

So in that simpler case, if  $h'' > 0$  at the critical point, then we lower the energy by sitting in the state  $|1\rangle$ , while if  $h'' < 0$  then we should sit in  $|0\rangle$ . This can be seen in the various examples that we explored in the previous section.

Now let's return to the multi-fermion case, with

$$H_{\text{Fermi}} = -\frac{1}{2}\partial_i\partial_j h[\psi^{\dagger i}, \psi^j]$$

At each critical point  $x = X$ , we should think of the Hessian  $\partial_i\partial_j h(X)$  as a matrix, with a collection of eigenvectors  $e_a^j$  and eigenvalues  $\lambda_a$ . In fact, to align with other conventions, it turns out to be best to think of the eigenvalue equation of the matrix  $-\partial_i\partial_j h$ ,

$$-(\partial_i\partial_j h) e_k^j = \lambda_k e_k^j \tag{1.27}$$

where  $k = 1, \dots, n$  labels the different eigenthings and shouldn't be summed over. The generalisation of the story above is now the following: for each negative eigenvalue  $\lambda_k < 0$ , we should excite the corresponding collection of fermions  $e_k^j \psi^{\dagger j}$ . Meanwhile, for each positive eigenvalue  $\lambda_k > 0$ , we should just leave well alone: we're better off in the unexcited state. At a given critical point  $x = X$ , the semi-classical ground state then sits in the part of the Hilbert space given by

$$|\text{ground}\rangle \sim \prod_{k \text{ with } \lambda_k < 0} (e_k^j \psi^{\dagger j})|0\rangle$$

We define the *Morse index* to be

$$\mu(X) = \text{The number of negative eigenvalues of } -\partial_i\partial_j h(X) \tag{1.28}$$

(We picked the eigenvalues of  $-\partial_i\partial_j h$  rather than  $+\partial_i\partial_j h$  so that this definition of the Morse index, in terms of negative rather than positive eigenvalues, is the standard one.) The ground state around the critical point  $X$  sits in the sector with  $\mu(X)$  excited fermions. In the geometrical language, this means that the ground state wavefunction is a  $p$ -form, where  $p = \mu(X)$  is the Morse index.

Now we can put everything together. We know that the Witten index only receives contributions from the ground states, and we now know that these are associated to critical points  $X$  of  $h$ , and live in the sector with  $\mu(X)$  excited fermions. We will assume that  $h(x)$  is chosen to be suitably generic so that there are no degenerate critical points. Then, using our previous result (1.24), we have

$$\text{Tr} (-1)^F e^{-\beta H} = \sum_X (-1)^{\mu(X)} \tag{1.29}$$

where the sum is over all critical points  $X$  of  $h$ .

Note that we're not assuming that all critical points of  $h$  correspond to true  $E = 0$  ground states of the theory. It may well be that some get lifted to non-zero energy and, later in these lectures, we'll put in some effort to understand when this happens. But that's not relevant for computing the Witten index since any such states must get lifted in pairs and so cancel out.

The same formula (1.29) also holds for our earlier model with a single  $x$  and  $\psi$ . There a maximum of  $h$  was necessarily followed by a minimum, so the sum over critical points could never exceed  $+1$  or drop below  $-1$ . Now, however, we could have multiple ground states. For example, we could have a situation where all the critical points  $X$  have  $\mu(X)$  even. In this case, they all contribute  $+1$  to the Witten index and each of them must correspond to a true,  $E = 0$  ground state of the system.

### 1.4.2 More Supersymmetry and Holomorphy

It is quite possible for a quantum system to be invariant under more than one supersymmetry transformation. The extended supersymmetry algebra replaces (1.1) with

$$\frac{1}{2}\{Q_\alpha, Q_\beta^\dagger\} = H\delta_{\alpha\beta} \quad \text{and} \quad \{Q_\alpha, Q_\beta\} = \{Q_\alpha^\dagger, Q_\beta^\dagger\} = 0 \quad (1.30)$$

with  $\alpha, \beta = 1, \dots, q$ . A Hamiltonian that can be written in this form is said to have  $N = 2q$  supersymmetries, with the 2 because each  $Q$  is complex. In this convention, the kind of quantum mechanics that we considered up until now is said to have  $N = 2$  supersymmetry. (I should warn you that the nomenclature for counting supersymmetry generators in quantum mechanics is not completely standard: things settle down once we go to higher dimensional quantum field theories.)

In this section, we'll construct a quantum mechanical model that has  $N = 4$  supersymmetry, meaning two complex supercharges  $Q_1$  and  $Q_2$  and their conjugates. Our strategy is to start with the Hamiltonian (1.25) with  $2n$  degrees of freedom. We'll split these up into two groups of  $n$ , related by supersymmetry as

$$\begin{aligned} x^i &\longleftrightarrow \psi_x^i \\ y^i &\longleftrightarrow \psi_y^i \end{aligned}$$

with  $i = 1, \dots, n$ . This supersymmetry is generated by the supercharge (1.26) which takes the form

$$\begin{aligned} Q_1 &= \left( p_{xi} - i \frac{\partial h}{\partial x^i} \right) \psi_x^i + \left( p_{yi} - i \frac{\partial h}{\partial y^i} \right) \psi_y^i \\ Q_1^\dagger &= \left( p_{xi} + i \frac{\partial h}{\partial x^i} \right) \psi_x^{\dagger i} + \left( p_{yi} + i \frac{\partial h}{\partial y^i} \right) \psi_y^{\dagger i} \end{aligned}$$

The supercharge depends on a single real function  $h(x, y)$ . The idea is to introduce a second supercharge that will relate the degrees of freedom in a different way, namely

$$\begin{aligned} x^i &\longleftrightarrow \psi_y^i \\ y^i &\longleftrightarrow \psi_x^i \end{aligned}$$

It takes some messing around to get the minus signs right, but it turns out that the following supercharge does the job

$$\begin{aligned} Q_2 &= \left( p_{xi} + i \frac{\partial h}{\partial x^i} \right) \psi_y^i - \left( p_{yi} + i \frac{\partial h}{\partial y^i} \right) \psi_x^i \\ Q_2^\dagger &= \left( p_{xi} - i \frac{\partial h}{\partial x^i} \right) \psi_y^{\dagger i} - \left( p_{yi} - i \frac{\partial h}{\partial y^i} \right) \psi_x^{\dagger i} \end{aligned}$$

The two supercharges  $Q_1$  and  $Q_2$  obey the algebra (1.30) but only if the function  $h(x, y)$  has some special properties. For example, we can compute the Hamiltonian in two different ways,

$$\begin{aligned} \{Q_1, Q_1^\dagger\} &= \sum_{i=1}^n \left( p_{xi}^2 + p_{yi}^2 + \left| \frac{\partial h}{\partial x^i} \right|^2 + \left| \frac{\partial h}{\partial y^i} \right|^2 \right) \\ &\quad - \sum_{i,j=1}^n \left( \frac{\partial^2 h}{\partial x^i \partial x^j} [\psi_x^{\dagger i}, \psi_x^j] + \frac{\partial^2 h}{\partial y^i \partial y^j} [\psi_y^{\dagger i}, \psi_y^j] + \frac{\partial^2 h}{\partial x^i \partial y^j} ([\psi_x^{\dagger i}, \psi_y^j] + [\psi_y^{\dagger j}, \psi_x^i]) \right) \end{aligned}$$

Alternatively, we have

$$\begin{aligned} \{Q_2, Q_2^\dagger\} &= \sum_{i=1}^n \left( p_{xi}^2 + p_{yi}^2 + \left| \frac{\partial h}{\partial x^i} \right|^2 + \left| \frac{\partial h}{\partial y^i} \right|^2 \right) \\ &\quad + \sum_{i,j=1}^n \left( \frac{\partial^2 h}{\partial x^i \partial x^j} [\psi_y^{\dagger i}, \psi_y^j] + \frac{\partial^2 h}{\partial y^i \partial y^j} [\psi_x^{\dagger i}, \psi_x^j] - \frac{\partial^2 h}{\partial x^i \partial y^j} ([\psi_y^{\dagger i}, \psi_x^j] + [\psi_x^{\dagger j}, \psi_y^i]) \right) \end{aligned}$$

The difference lies in the second line, where the  $\psi_x$  and  $\psi_y$  fermions are exchanged, together with some minus signs. At first glance, it looks like these are simply different Hamiltonians. However, all is not lost: these two Hamiltonians coincide if the function  $h(x, y)$  obeys

$$\frac{\partial^2 h}{\partial x^i \partial x^j} = - \frac{\partial^2 h}{\partial y^i \partial y^j} \quad \text{and} \quad \frac{\partial^2 h}{\partial x^i \partial y^j} = \frac{\partial^2 h}{\partial y^i \partial x^j} \quad (1.31)$$

There's a much nicer way of writing these conditions: as we will now see, they are telling us that  $h(x, y)$  is related to a holomorphic function.

## Complex Variables

We introduce the complex coordinates

$$z^i = x^i + iy^i \quad \text{and} \quad \bar{z}^i = x^i - iy^i$$

Notice the extra bar on the  $\bar{i} = 1, \dots, n$  index on the conjugate  $z^\dagger = \bar{z}$ ; it's a fairly common notation when dealing with complex coordinates. The corresponding derivative operators are

$$\partial_i = \frac{1}{2} \left( \frac{\partial}{\partial x^i} - i \frac{\partial}{\partial y^i} \right) \quad \text{and} \quad \bar{\partial}_i = \frac{1}{2} \left( \frac{\partial}{\partial x^i} + i \frac{\partial}{\partial y^i} \right)$$

which obeys  $\partial_i z^j = \delta_i^j$  and  $\bar{\partial}_i \bar{z}^j = \delta_i^j$  and  $\partial_i \bar{z}^j = \bar{\partial}_i z^j = 0$ .

Now consider a holomorphic function  $W(z)$  which depends only on the  $z^i$  and not on  $\bar{z}^i$ . If we decompose this in terms of a real and imaginary piece

$$W(z) = -h(x, y) - ig(x, y) \tag{1.32}$$

then the Cauchy-Riemann equations read

$$\frac{\partial W}{\partial \bar{z}^i} = 0 \quad \Rightarrow \quad \frac{\partial h}{\partial x^i} = \frac{\partial g}{\partial y^i} \quad \text{and} \quad \frac{\partial h}{\partial y^i} = -\frac{\partial g}{\partial x^i}$$

It is simple to show that these then imply the requirements (1.31).

This motivates us to frame the theory with  $N = 4$  supersymmetry in terms of complex variables rather than real variables. In addition to the complex coordinates  $z^i$ , we also introduce complex momenta

$$p_i = \frac{1}{2} (p_{xi} - ip_{yi}) \quad \text{and} \quad \bar{p}_i = \frac{1}{2} (p_{xi} + ip_{yi})$$

as well as “complex” Grassmann variables. Here the word “complex” is in inverted commas because our original Grassmann variables were already complex; we just introduce different linear combinations

$$\begin{aligned} \Psi^i &= \psi_x^i + i\psi_y^i & \text{and} & \quad \bar{\Psi}^i = \psi_x^{\dagger i} - i\psi_y^{\dagger i} \\ \tilde{\Psi}^i &= \psi_x^{\dagger i} + i\psi_y^{\dagger i} & \text{and} & \quad \tilde{\bar{\Psi}}^i = \psi_x^i - i\psi_y^i \end{aligned}$$

We've now abandoned the  $\dagger$  notation for complex conjugation and resorted instead to the barred notation. (If nothing else, it is easier to write bars when doing long

calculations.) Finally, we have combinations of supercharges

$$\begin{aligned}
Q_+ &= \frac{1}{2}(Q_1 + iQ_2) = p_i \Psi^i + \frac{i}{2} \bar{\partial}_i \bar{W} \bar{\Psi}^{\bar{i}} \\
Q_+^\dagger &= \frac{1}{2}(Q_1^\dagger - iQ_2^\dagger) = \bar{p}_i \bar{\Psi}^{\bar{i}} - \frac{i}{2} \partial_i W \tilde{\Psi}^i \\
Q_- &= \frac{1}{2}(Q_1 - iQ_2) = \bar{p}_i \bar{\Psi}^{\bar{i}} + \frac{i}{2} \partial_i W \Psi^i \\
Q_-^\dagger &= \frac{1}{2}(Q_1^\dagger + iQ_2^\dagger) = p_i \tilde{\Psi}^i - \frac{i}{2} \partial_i \bar{W} \bar{\Psi}^{\bar{i}}
\end{aligned}$$

These obey the extended supersymmetry algebra  $\{Q_\alpha, Q_\beta^\dagger\} = H\delta_{\alpha,\beta}$ , now with  $\alpha, \beta = +, -$ .

The flurry of complexified definitions conspire to make the theory look somewhat simpler. In the Lagrangian picture, it takes the form

$$\begin{aligned}
L = \sum_{i=1}^n \left( |\dot{z}_i|^2 + i\tilde{\Psi}^i \partial_t \bar{\Psi}^{\bar{i}} + i\bar{\Psi}^{\bar{i}} \partial_t \Psi^i - \frac{1}{4} |\partial_i W|^2 \right) \\
- \frac{1}{2} \sum_{i,j} \left( \partial_i \partial_j W \Psi^i \tilde{\Psi}^j + \bar{\partial}_i \bar{\partial}_j \bar{W} \bar{\Psi}^{\bar{i}} \bar{\Psi}^{\bar{j}} \right) \quad (1.33)
\end{aligned}$$

Supersymmetric Lagrangians of this kind, involving complex scalar fields and fermions, are usually referred to as *Landau-Ginzburg* theories. This is a nod to the Landau-Ginzburg theories that we met when discussing phase transitions in [Statistical Physics](#). But it's not a very good nod. In particular, the theory (1.25) with just a single supersymmetry is just as much related to the kinds of models that Landau and Ginzburg considered but is never given this name in the context of supersymmetry. It's best to think of the name "Landau-Ginzburg" for the Lagrangian (1.33) as merely a quirk of history and forget that the term is also used elsewhere in physics.

The Landau-Ginzburg Lagrangian depends on a single holomorphic function  $W(z)$ . This is known as the *superpotential*. The fact that extended supersymmetry comes hand in hand with holomorphy and associated ideas in complex analysis is extremely important. We will not discuss quantum mechanics with  $N = 4$  supersymmetry in these lectures, but it's not for want of interesting content. In particular, there is a beautiful relationship to a form of complex geometry known as "Kähler geometry" that underlies many of the most interesting results in this subject.

Furthermore, when we go to higher dimensional field theories, supersymmetry generators are associated to spinors and these necessarily have more than one component.

This means that in, for example,  $d = 3 + 1$  dimensions, the simplest supersymmetric theories have the form (1.33) and are based on complex, rather than real variables. In that context, the holomorphy of the superpotential goes a long way towards allowing us to solve some complicated features of supersymmetric quantum field theories. This is covered in some detail in the lectures on [Supersymmetric Field Theory](#).

## The Ground States

Finally, we can turn to some physics of the theory (1.33). As previously, we can ask how many ground states the theory has. The semi-classical ground states are associated to critical points of the superpotential,

$$\partial_i W = 0 \quad \text{for all } i = 1, \dots, n$$

We know from our discussion in Section 1.4.1 what we should do next: we compute the Morse index for each critical point, meaning the number of positive eigenvalues of the Hessian of  $h$ . But this is trivial for a holomorphic function  $W(z)$ . For example, if there is a critical point near the origin, we can expand (after a suitable diagonalisation)

$$W(z) \approx \sum_i \lambda_i (z^i)^2 + \dots$$

In terms of our real variables,  $(z^i)^2 = (x^i)^2 - (y^i)^2 + 2ix^iy^i$ , while our original function  $h(x, y)$  is given, from (1.32), as

$$h(x, y) = -\text{Re } W = -\sum_i \lambda_i \left( (x^i)^2 - (y^i)^2 \right) + \dots$$

We learn that, because of the holomorphy of  $W$ , for every positive eigenvalue of the Hessian of  $h$ , there is a corresponding negative eigenvalue. This ensures that every critical point has morse index (1.28) given by  $n$  and each contributes exactly the same to the Witten index (1.29) which becomes

$$\text{Tr} (-1)^F e^{-\beta H} = (-1)^n \times \text{Number of critical points of } W$$

We learn that in theories with  $N = 4$  supersymmetry, every critical point of  $W$  is a true  $E = 0$  ground state of the quantum theory.

### 1.4.3 Less Supersymmetry and Spinors

It's also possible to consider theories with less supersymmetry than our starting point. In fact, this is easy to achieve. We return to our theory with  $N = 2$  supersymmetry and impose a reality condition on the Grassmann variables

$$\psi^{\dagger i} = \psi^i$$

Real quantum mechanical Grassmann variables like this are called *Majorana modes* or *Majorana fermions*.

For our current purposes, it will suffice to discuss just the free theory,

$$S = \int dt \sum_{i=1}^n \left( \frac{1}{2} \dot{x}^i \dot{x}^i + \frac{i}{2} \psi^i \dot{\psi}^i \right) \quad (1.34)$$

This is invariant under a single real supercharge,

$$Q = \sum_i \dot{x}^i \psi^i$$

which obeys  $Q^\dagger = Q$  and generates the supersymmetry transformation

$$\delta x^i = \epsilon \psi^i \quad \text{and} \quad \delta \psi^i = i \epsilon \dot{x}^i$$

This is usually referred to as  $N = 1$  supersymmetry. (You will sometimes see the terminology  $N = \frac{1}{2}$  supersymmetry in the literature, counting complex supercharges rather than real.)

Here our interest lies in a very specific property of these theories: how should we think of the internal degrees of freedom generated by the real fermions  $\psi^i$ ? There is a very pretty answer to this question. To see this, first note that the momentum conjugate to the fermion is

$$\frac{\partial L}{\partial \dot{\psi}^i} = -\frac{i}{2} \psi^i$$

The canonical commutation relation for the fermion is then

$$\{\psi^i, \psi^j\} = 2\delta^{ij} \quad i, j = 1, \dots, n$$

But this is a very familiar equation: it is the Clifford algebra, usually written in terms of gamma matrices

$$\{\gamma^i, \gamma^j\} = 2\delta^{ij} \quad i, j = 1, \dots, n$$

This means that the fermions in this theory should be viewed as gamma matrices! The Clifford algebra has a unique irreducible representation of dimension  $2^{n/2}$  if  $n$  is even and  $2^{(n-1)/2}$  if  $n$  is odd. This strongly suggests that the internal degrees of freedom of the particle described by the action (1.34) have something to do with spinors on  $\mathbb{R}^n$ .

It is straightforward to construct these internal degrees of freedom. First, let's assume that  $n$  is even. (We will discuss the case of  $n$  odd below.) We pair up the Majorana modes into complex fermions

$$c^i = \frac{1}{2} (\psi^{2i} + i\psi^{2i-1}) \quad i = 1, \dots, \frac{n}{2}$$

Then the complex  $c^i$  operators form the usual algebra of fermionic creation and annihilation operators that we're used to

$$\{c^{\dagger i}, c^j\} = \delta^{ij} \quad \text{and} \quad \{c^{\dagger i}, c^{\dagger j}\} = \{c^i, c^j\} = 0$$

and we can use them to build the familiar fermionic Fock space starting with  $|0\rangle$  that obeys  $c^i|0\rangle = 0$  and then acting with  $c^{\dagger i}$ . Following the discussion in Section 1.4.1, we see that the fermions fill out an internal space with

$$\text{Dimension of Internal Space} = 2^{n/2}$$

This is precisely the dimension of a Dirac spinor on  $\mathbb{R}^n$ .

There is more to say about these spinors. Under a rotation in  $\mathbb{R}^n$ , the Dirac spinor transforms in the representation generated by  $\Sigma^{ij} = \frac{1}{4}[\gamma^i, \gamma^j]$ . (See the lectures on [Quantum Field Theory](#) for more details of this.) However, in even dimension, as we have here, this is not an irreducible representation. It is composed of two smaller representations known as *chiral spinors* or *Weyl spinors*.

These arise because we can always construct an operator  $\hat{\gamma}$  that is analogous to  $\gamma^5$  in four dimensions. In general, this is given by multiplying all the gamma matrices together, with a suitable factor of  $i$  to ensure Hermiticity,

$$\hat{\gamma} = i^{n/2} \gamma^1 \dots \gamma^n$$

This obeys  $\hat{\gamma}^2 = 1$  and  $\{\hat{\gamma}, \gamma^i\} = 0$ . The existence of the  $\hat{\gamma}$  operator means that all the internal states can be decomposed into two different camps: those with eigenvalue  $\hat{\gamma} = +1$  and those with eigenvalue  $\hat{\gamma} = -1$ . In the language of the Dirac equation, these are spinors of different chirality.

In the context of our supersymmetric quantum mechanics, this  $\hat{\gamma}$  operator has a very natural meaning. The eigenvalues are simply states with an even or odd number of  $c^\dagger$  operators excited. In other words, this plays the role of our fermion number.

$$\hat{\gamma} = (-1)^F$$

This means that  $\hat{\gamma}$  determines whether states live in  $\mathcal{H}_B$  or  $\mathcal{H}_F$ .

The punchline of this argument is that quantising real fermions, appropriate for  $N = 1$  supersymmetry, gives Dirac spinors on  $\mathbb{R}^n$ , at least for  $n$  even. These have dimension  $2^{n/2}$ . Meanwhile, while quantising complex fermions, appropriate for  $N = 2$  supersymmetry, gives forms on  $\mathbb{R}^n$ . These have dimension  $2^n$ . We'll have use for quantum mechanics with  $N = 1$  supersymmetry in Section 3.3 where we discuss the Atiyah-Singer index theorem.

As an aside, clearly the construction of spinors and forms on  $\mathbb{R}^n$  from Grassmann degrees of freedom is closely related. This also suggests that you can take  $2^{n/2}$  different Dirac spinors and bundle them together to look like forms. Such a construction is called *Kähler-Dirac fermions*. It won't play a role in these lectures, but arises in a number of other areas of physics including topological twisting of field theories and lattice gauge theory where it goes by the name of *staggered fermions*.

### The Case of $n$ Odd: A Subtle Anomaly

We still have to understand the case of  $n$  odd. Here there is a surprise. Quantum mechanical theories with an odd number of Majorana modes don't make any sense! They are an example of what is sometimes called an *anomalous* quantum theory: a seemingly sensible classical theory that cannot be quantised.

The argument is straightforward. Consider two, non-interacting Majorana fermions,  $\psi^1$  and  $\psi^2$ . From the discussion above, we can construct a single complex fermion  $c = (\psi^1 + i\psi^2)/\sqrt{2}$  and this acts on a two-dimensional Hilbert space spanned by  $|0\rangle$  and  $c^\dagger|0\rangle$ .

But, by the factorisation of Hilbert spaces, that means that a single Majorana fermion, say  $\psi^1$ , must act on a Hilbert space of dimension  $\sqrt{2}$ . And that's nonsense! You can reach the same conclusion if you use the path integral to compute  $\text{Tr} e^{-\beta H}$ , which just counts the dimension of the Hilbert space when  $H = 0$ . Again, after suitable regularisation, you find  $\sqrt{2}$ .

For us, this means that theories with  $N = 1$  supersymmetry are restricted to describe a particle moving in an even dimensional space, like  $\mathbb{R}^n$  with  $n$  even.

## 2 Supersymmetry and the Path Integral

In addition to making the symmetry aspect of supersymmetry manifest, the Lagrangian description of the quantum mechanics has one additional advantage: it allows us to bring the path integral to bear on the problem.

We'll make plenty of use of the path integral in later studies of supersymmetric systems. The purpose of this section is to understand some of the basic properties of the quantum mechanical path integral and how we can use it to compute quantities of interest in supersymmetric theories.

### 2.1 The Partition Function and the Index

Let's start with a purely bosonic system, with the familiar action

$$S = \int dt \frac{1}{2} \dot{x}^2 - V(x) \quad (2.1)$$

In statistical mechanics, we typically want to compute the partition function

$$Z = \text{Tr} e^{-\beta H}$$

How do we compute this using path integrals?

Our starting point is Feynman's expression for the propagator. Take a particle that sits at point  $x_i$  at time  $t_i$ . The quantum amplitude for it to be at point  $x_f$  at a later time  $t_f$  has the beautiful path integral expression

$$\langle x_f | e^{-iH(t_f - t_i)} | x_i \rangle = \int_{x(t_i) = x_i}^{x(t_f) = x_f} \mathcal{D}x(t) e^{iS[x(t)]} \quad (2.2)$$

Note that the initial and final states  $|x_i\rangle$  and  $|x_f\rangle$  appear on the right-hand-side as boundary conditions for the paths: we integrate over all paths  $x(t)$  such that  $x(t_i) = x_i$  and  $x(t_f) = x_f$ .

Our goal now is to manipulate (2.2) so that the left-hand-side looks like the partition function  $Z$ . There are a number of differences that we need to fix. First, the time evolution operator in quantum mechanics is unitary,  $e^{-iHt}$ . Meanwhile, in statistical mechanics the relevant operator is  $e^{-\beta H}$ , with a minus sign in the exponent rather than a factor of  $i$ . To deal with this, we work in *imaginary time*,

$$\tau = it$$

So  $e^{-iHt} = e^{-H\tau}$ . On the right-hand-side, we write the action in Euclidean time so it becomes

$$S = +i \int d\tau \frac{1}{2} \left( \frac{dx}{d\tau} \right)^2 + V(x) = iS_E$$

Here  $S_E$  is the *Euclidean action*. In quantum field theory, this operation is tantamount to Wick rotation. The Feynman expression (2.2) then becomes

$$\langle x_f | e^{-H(\tau_f - \tau_i)} | x_i \rangle = \int_{x(\tau_i)=x_i}^{x(\tau_f)=x_f} \mathcal{D}x(\tau) e^{-S_E[x(\tau)]} \quad (2.3)$$

That's fixed up the minus signs and factors of  $i$ . Next up is the trace in the partition function. This means that we sum over a basis of states. If we choose that basis to be position eigenstates, then we have

$$Z = \text{Tr} e^{-\beta H} = \int_{-\infty}^{+\infty} dx \langle x | e^{-\beta H} | x \rangle$$

Comparing this to (2.3), we have

$$Z = \int_{-\infty}^{+\infty} dx \langle x | e^{-\beta H} | x \rangle = \int_{-\infty}^{+\infty} dx_i \int_{x(\tau_i)=x_i}^{x(\tau_f)=x_i} \mathcal{D}x(\tau) e^{-S_E[x(\tau)]}$$

where  $\tau_f = \tau_i + \beta$  and the important point is that we now integrate over paths where the particle comes back to where it started:  $x_f = x_i$ . Furthermore, we integrate over all possible starting points  $x_i$ . This gives our final expression for the partition function which we write as

$$Z = \text{Tr} e^{-\beta H} = \int_{x(\beta)=x(0)} \mathcal{D}x(\tau) e^{-S_E[x(\tau)]}$$

where now the boundary conditions just tell us that we should integrate over all possible closed paths. Equivalently, we can implement this condition by insisting that we work in periodic Euclidean time, so that  $\tau$  is a coordinate on a circle  $\mathbf{S}^1$ , with

$$\tau \equiv \tau + \beta$$

Although we've derived this punchline in the context of quantum mechanics, it also works in quantum field theory. If you want to compute the thermal partition function of any quantum field theory, you simply need to work in periodic, Euclidean time. This will tell you information about the equilibrium properties of the system at temperature  $T = 1/\beta$ .

### 2.1.1 An Example: The Harmonic Oscillator

To get a sense for how these calculations work, let's look at everyone's favourite example: the harmonic oscillator. This, of course, takes the form (2.1) with the potential

$$V(x) = \frac{1}{2}\omega^2 x^2$$

We know that the harmonic oscillator has energy levels  $E = \omega(n + \frac{1}{2})$  with  $n = 0, 1, \dots$  (assuming that  $\omega > 0$ ). This means that, in this case, we can just compute the partition function by summing over all states.

$$Z = e^{-\beta\omega/2} \sum_{n=0}^{\infty} e^{-n\beta\omega} = \frac{e^{-\beta\omega/2}}{1 - e^{-\beta\omega}} \quad (2.4)$$

How does the path integral reproduce this? We have

$$Z = \int \mathcal{D}x(\tau) \exp\left(-\frac{1}{2} \int d\tau x \left(-\frac{d^2}{d\tau^2} + \omega^2\right) x\right)$$

where we've left the periodic boundary conditions implicit and integrated by parts in the Euclidean action to highlight the fact that the resulting path integral takes a Gaussian form. If we extrapolate from finite-dimensional Gaussian integrals, we find ourselves with the slightly formal expression

$$Z = \det\left(-\frac{d^2}{d\tau^2} + \omega^2\right)^{-1/2}$$

We should think of this determinant as the product of eigenvalues of the differential operator. The eigenfunctions of this operator are

$$x(\tau) = e^{ik\tau} \quad \Rightarrow \quad \left(-\frac{d^2}{d\tau^2} + \omega^2\right)x(\tau) = (k^2 + \omega^2)x(\tau)$$

but we should remember that we're working on a circle with periodic boundary conditions so we require  $x(\tau + \beta) = x(\tau)$ . This restricts the  $k$  values to be quantised

$$k = \frac{2\pi n}{\beta} \quad n \in \mathbb{Z}$$

This, of course, is just the usual calculations that we do in our first course on [Quantum Mechanics](#). The novelty here is that we now need to multiply all these eigenvalues together

$$Z = \prod_{n=-\infty}^{+\infty} \left[ \left(\frac{2\pi n}{\beta}\right)^2 + \omega^2 \right]^{-1/2} = \frac{1}{\omega} \prod_{n=1}^{\infty} \left[ \left(\frac{2\pi n}{\beta}\right)^2 + \omega^2 \right]^{-1}$$

In the second equality we've taken out the  $n = 0$  term, and then used the fact that  $\pm n$  give the same contribution to remove the square-root factor at the expense of restricting the product to positive integers. We can rewrite the resulting expression as a product of two terms, each itself an infinite product

$$Z = \frac{1}{\omega} \prod_{n'=1}^{\infty} \left( \frac{2\pi n'}{\beta} \right)^{-2} \prod_{n=1}^{\infty} \left[ 1 + \left( \frac{\beta\omega}{2\pi n} \right)^2 \right]^{-1} \quad (2.5)$$

The second of these expressions is convergent and the result is well known:

$$\prod_{n=1}^{\infty} \left[ 1 + \left( \frac{\beta\omega}{2\pi n} \right)^2 \right] = \frac{2}{\beta\omega} \sinh \frac{\beta\omega}{2} \quad (2.6)$$

We won't prove this result here, but just note that it follows immediately from Euler's product formula for sine,

$$\sin(\pi z) = \pi z \prod_{n=1}^{\infty} \left( 1 - \frac{z^2}{n^2} \right)$$

As an obvious sanity check, it's clear that both sides of this equation have the same zeros. A full proof of the equivalence is not too hard, but a little involved.

That leaves us with the first infinite product in (2.5) to deal with. And that's more tricky because it diverges. To better understand such terms, we should really go back and dissect the path integral to figure out where it came from. (For example, the partition function should be dimensionless but this term has dimension of  $[\text{Energy}]^{2\infty}$  which is a hint that we didn't define our measure very well.) However, in the spirit of this course we're going to treat this term as blithely as possible. And, for those physicists of a blithe disposition, there are few tools more useful than zeta function regularisation.

The zeta function is defined, for  $\text{Re}(s) > 1$ , by the sum

$$\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$$

However,  $\zeta(s)$  is defined for all values of  $s$ . The idea is that we use this to give meaning to divergent sums. For example, we could think of the sum of all positive integers as  $\zeta(-1) = -1/12$ . Although these zeta function games seem rather inane when you first meet them, the magic is that they tend to give the right answers when used to regulate divergences in quantum field theory. (For example, in the lectures on [String Theory](#) we first invoked the unconvincing  $\zeta(-1) = -1/12$  argument to compute the critical dimension of the string, and then spent a significant amount of time rederiving this using conformal field theory techniques where the divergences were absent.)

Let's see how we can put the zeta function to work for the harmonic oscillator. We first introduce the related function

$$\zeta_1(s) = \left(\frac{\beta}{2\pi}\right)^{2s} \zeta(2s) = \sum_{n=1}^{\infty} \left(\frac{2\pi n}{\beta}\right)^{-2s}$$

Now we take the derivative with respect to  $s$ . Note that, if  $y = a^x$  then  $dy/dx = y \log a$ . So

$$\begin{aligned} \zeta_1'(s) &= 2 \left(\frac{\beta}{2\pi}\right)^{2s} \left( \log\left(\frac{\beta}{2\pi}\right) \zeta(2s) + \zeta'(2s) \right) \\ &= \sum_{n=1}^{\infty} \left(\frac{2\pi n}{\beta}\right)^{-2s} \log\left(\frac{2\pi n}{\beta}\right)^{-2} \end{aligned}$$

Evaluated at  $s = 0$ , we have

$$\zeta_1'(0) = 2 \log\left(\frac{\beta}{2\pi}\right) \zeta(0) + 2\zeta'(0) = \sum_{n=1}^{\infty} \log\left(\frac{2\pi n}{\beta}\right)^{-2}$$

Or, taking the exponential of both sides,

$$\prod_{n=1}^{\infty} \left(\frac{2\pi n}{\beta}\right)^{-2} = \left(\frac{\beta}{2\pi}\right)^{2\zeta(0)} e^{2\zeta'(0)}$$

Now we need the values  $\zeta(0) = -\frac{1}{2}$  and  $\zeta'(0) = -\frac{1}{2} \log(2\pi)$ . Combining these gives the zeta-tamed value for the divergent product

$$\prod_{n=1}^{\infty} \left(\frac{2\pi n}{\beta}\right)^{-2} = \frac{1}{\beta} \tag{2.7}$$

We can see immediately that, despite the dubious route to get there, the end result is plausible. This is because the  $1/\beta$  factor from the regularised product combines with the  $1/\omega$  factor that sits in front of (2.5) to ensure that the partition function is dimensionless, as it should be.

Putting this together with our convergent product (2.6), we get our end result for the path integral

$$Z = \frac{1}{2 \sinh(\beta\omega/2)} \tag{2.8}$$

But this is what we want: it agrees with the harmonic oscillator partition function (2.4) computed through more traditional means.

One last, trivial observation before we move on. The harmonic oscillator potential depends, of course, on  $\omega^2$ . In the derivation above, we assumed that  $\omega > 0$ . In what follows, a better answer for the partition function is

$$Z = \frac{1}{2 \sinh(\beta|\omega|/2)} \tag{2.9}$$

This trivial amendment will be important to remember later when we discuss the supersymmetric oscillator.

### 2.1.2 Fermions: Periodic or Anti-Periodic?

When dealing with supersymmetric systems, our theory necessarily contains fermionic, or Grassmann, variables. And these bring a new subtlety to the problem.

In quantum field theories in higher dimensions, fermions famously come with a minus sign issue: rotate a fermionic field by  $2\pi$  and it doesn't come back to itself, but picks up a minus sign. This same minus sign manifests itself when computing the thermal partition function.

As we saw in Section 2.1, if we want to compute  $Z = \text{Tr} e^{-\beta H}$  then we should work in Euclidean time with period  $\beta$ . The bosonic fields  $x(t)$  are given periodic boundary conditions

$$x(\tau + \beta) = x(\tau)$$

But for the fermionic fields, that minus sign suggest two possibilities: we could have periodic or anti-periodic boundary conditions

$$\psi(\tau + \beta) = \psi(\tau) \quad \text{or} \quad \psi(\tau + \beta) = -\psi(\tau)$$

Relatedly, there are two natural partition functions that we could construct for fermions. In addition to the thermal partition function  $\text{Tr} e^{-\beta H}$ , we could also consider the quantity  $\text{Tr}(-1)^F e^{-\beta H}$ . In supersymmetric quantum mechanics,  $\text{Tr}(-1)^F e^{-\beta H}$  is the Witten index and is necessarily an integer. But, for a general fermionic system it is just a different way to sum the states, weighted by an extra minus sign. I'll refer to the quantity  $\text{Tr}(-1)^F e^{-\beta H}$  as an "index" in both supersymmetric and non-supersymmetric theories, although strictly this terminology should be reserved for the former case.

It seems plausible that inserting a factor of  $(-1)^F$  in the trace would flip the sign of the fermion as we go around the Euclidean temporal circle. But which boundary condition corresponds to the thermal partition function, and which to the index?

As always, the right answer can be found by going back to first principles and looking at how one constructs the path integral from small, but finite, time steps. Here, however, we will simply give the answer and then provide some motivation. The answer is that the thermal partition function requires *anti-periodic* boundary conditions for fermions,

$$\text{Anti-Periodic: } Z = \text{Tr} e^{-\beta H} = \int_{\psi(\beta)=-\psi(0)} \mathcal{D}\psi^\dagger \mathcal{D}\psi e^{-S_E[\psi, \psi^\dagger]}$$

Meanwhile, the index is computed with periodic boundary conditions:

$$\text{Periodic: } \text{Tr} (-1)^F e^{-\beta H} = \int_{\psi(\beta)=\psi(0)} \mathcal{D}\psi^\dagger \mathcal{D}\psi e^{-S_E[\psi, \psi^\dagger]}$$

To motivate this result, we will calculate the path integral in a particularly simple case.

### The Fermionic Oscillator

The simple model that we'll use as a testing ground is a free fermion with action

$$S = \int dt \left[ i\psi^\dagger \dot{\psi} - \omega \psi^\dagger \psi \right]$$

This is nothing complicated: it is the Lagrangian description for a two state system. As we've seen previously, the canonical commutation relations are  $\{\psi, \psi\} = \{\psi^\dagger, \psi^\dagger\} = 0$  and  $\{\psi, \psi^\dagger\} = 1$  and these naturally act on a two-dimensional Hilbert space spanned by  $|0\rangle$  and  $|1\rangle$  such that

$$\psi|0\rangle = 0 \quad \text{and} \quad |1\rangle = \psi^\dagger|0\rangle$$

The Hamiltonian of this system is

$$H = \frac{1}{2}\omega[\psi^\dagger, \psi] \quad \Rightarrow \quad H|0\rangle = -\frac{\omega}{2}|0\rangle \quad \text{and} \quad H|1\rangle = +\frac{\omega}{2}|1\rangle$$

Note that we've chosen the symmetric operator ordering for the Hamiltonian, so that the energies are  $E = \pm\omega/2$ . In the absence of supersymmetry, there is nothing that enforces this upon us and other orderings will give energies shifted by  $E + \text{constant}$ . However, we will see below that the naive implementation of the path integral also gives this symmetric choice of energies.

For the two state system, the computation of the thermal partition function using the Hamiltonian approach is a trivial calculation: we get

$$Z = \text{Tr} e^{-\beta H} = e^{-\beta\omega/2} + e^{+\beta\omega/2} \quad (2.10)$$

We define the fermion number  $F = \psi^\dagger \psi$ , so  $F|0\rangle = 0$  and  $F|1\rangle = |1\rangle$ . Then the index differs from the partition function just by a minus sign

$$\text{Tr} (-1)^F e^{-\beta H} = -e^{-\beta\omega/2} + e^{+\beta\omega/2} \quad (2.11)$$

Clearly the index *isn't* independent of  $\beta$  for this simple model: that is only true for supersymmetric systems.

Our challenge is to reproduce these two results from the path integral and use this to confirm which boundary condition gives which quantity. For both choices of boundary condition, the starting point is the same: the Euclidean action is

$$S_E[\psi^\dagger, \psi] = \int d\tau \left[ \psi^\dagger \frac{d\psi}{d\tau} + \omega \psi^\dagger \psi \right]$$

The fermionic path integral is Gaussian. By dint of the complex Grassmann nature of the integration variables, we get  $\det$  rather than  $\det^{-1/2}$ , so that

$$\int \mathcal{D}\psi^\dagger \mathcal{D}\psi e^{-S_E[\psi^\dagger, \psi]} = \det \left( \frac{d}{d\tau} + \omega \right)$$

We again think of the determinant as the product of eigenvalues. The eigenfunctions have the same form as before

$$\psi(\tau) = \eta_0 e^{ik\tau} \quad \Rightarrow \quad \left( \frac{d}{d\tau} + \omega \right) \psi = (ik + \omega)\psi$$

for some constant Grassmann parameter  $\eta_0$ . The difference between periodic and anti-periodic boundary conditions comes in the allowed values of  $k$ . We have

$$\begin{aligned} \text{Periodic : } \psi(\tau + \beta) = \psi(\tau) &\quad \Rightarrow \quad k = \frac{2\pi n}{\beta} \\ \text{Anti-Periodic : } \psi(\tau + \beta) = -\psi(\tau) &\quad \Rightarrow \quad k = \frac{2\pi(n - 1/2)}{\beta} \end{aligned}$$

with  $n \in \mathbb{Z}$ . We see that the modes are, up to a normalisation, either integer or half-integer valued depending on the choice of boundary conditions.

Let's start with the periodic case. We have

$$\begin{aligned}
\text{Periodic: } \det\left(\frac{d}{d\tau} + \omega\right) &= \prod_{n \in \mathbb{Z}} \left(\frac{2\pi i n}{\beta} + \omega\right) \\
&= \omega \prod_{n=1}^{\infty} \left(\left(\frac{2\pi n}{\beta}\right)^2 + \omega^2\right) \\
&= \omega \prod_{n'=1}^{\infty} \left(\frac{2\pi n'}{\beta}\right)^2 \prod_{n=1}^{\infty} \left(1 + \left(\frac{\beta\omega}{2\pi n}\right)^2\right) \\
&= 2 \sinh\left(\frac{\beta\omega}{2}\right) \tag{2.12}
\end{aligned}$$

where, in the last line, we've used our previous expressions for the convergent product (2.6) and the divergent product, tamed by zeta function regularisation (2.7). As promised, this coincides with the index  $\text{Tr}(-1)^F e^{-\beta H}$  that we computed in (2.11). (Actually it differs by a minus sign, but this is simply the convention for  $F$ .)

Meanwhile, with anti-periodic boundary conditions, we have

$$\text{Anti-Periodic: } \det\left(\frac{d}{d\tau} + \omega\right) = \prod_{n \in \mathbb{Z}} \left(\frac{2\pi i(n - 1/2)}{\beta} + \omega\right)$$

The modes  $k$  come in  $\pm$  pairs with  $n$  pairing with  $-n + 1$ . (So, for example,  $n = 1$  pairs up with  $n = 0$  since both have  $k = \pm\pi/\beta$ .) We use this to rewrite the product as

$$\begin{aligned}
\det\left(+\frac{d}{d\tau} + \omega\right) &= \prod_{n=1}^{\infty} \left(\left(\frac{2\pi(n - 1/2)}{\beta}\right)^2 + \omega^2\right) \\
&= \prod_{n'=1}^{\infty} \left(\frac{2\pi(n' - 1/2)}{\beta}\right)^2 \prod_{n=1}^{\infty} \left(1 + \left(\frac{\beta\omega}{2\pi(n - 1/2)}\right)^2\right)
\end{aligned}$$

Again, the determinant factorises into two inner products. Again, the second of these is convergent and has a well known form (that, once more, we won't prove),

$$\prod_{n=1}^{\infty} \left(1 + \left(\frac{\beta\omega}{2\pi(n - 1/2)}\right)^2\right) = \cosh\left(\frac{\beta\omega}{2}\right)$$

We're left, however, with the first infinite product and this is clearly divergent. As before, we turn to zeta function regularisation for refuge. The same argument that we used for the bosonic oscillator can be invoked here too, now applied to the so-called Hurwitz zeta function

$$\zeta(s, 1/2) = \sum_{n=0}^{\infty} \frac{1}{(n + 1/2)^s}$$

The upshot is that, with anti-periodic boundary conditions, the path integral gives

$$\det \left( \frac{d}{d\tau} + \omega \right) \Big|_{\text{anti-periodic}} = 2 \cosh \left( \frac{\beta\omega}{2} \right)$$

This reproduces the thermal partition function (2.10).

It will not have escaped your attention that the path integral calculation was a lot of work to get the partition function for a two state system. However, as we come to consider more complicated quantum mechanical models, including higher dimensional field theories, the path integral starts to come into its own and, ultimately, is much more convenient than canonical quantisation.

### 2.1.3 The Witten Index Revisited

It's useful to understand why, from the path integral perspective, the Witten index is always an integer for supersymmetric theories. After all, something magical must happen where we do an infinite dimensional integral but, regardless of the parameters in the integrand, we always get an integer. How does this come about? The answer is a rather special property of supersymmetric path integrals known as *localisation*.

To see how this works, we'll revert to the simplest system of a particle with spin on a line. In Euclidean time, the action (1.14) becomes

$$S_E[x, \psi, \psi^\dagger] = \oint d\tau \left[ \frac{1}{2} \left( \frac{dx}{d\tau} \right)^2 + \psi^\dagger \frac{d\psi}{d\tau} + \frac{1}{2} h'^2 - h'' \psi^\dagger \psi \right] \quad (2.13)$$

where the  $\oint$  is there to remind us that we're working in periodic time. The Euclidean action is invariant under the Wick rotated supersymmetry transformations (1.19), which read

$$\delta x = \epsilon^\dagger \psi - \epsilon \psi^\dagger \quad , \quad \delta \psi = \epsilon \left( \frac{dx}{d\tau} + h' \right) \quad , \quad \delta \psi^\dagger = \epsilon^\dagger \left( -\frac{dx}{d\tau} + h' \right) \quad (2.14)$$

The bosonic field  $x(\tau)$  is always periodic:  $x(\tau) = x(\tau + \beta)$ . But that means that the supersymmetry transformations (2.14) only hold if  $\psi$  is also periodic:  $\psi(\tau) = \psi(\tau + \beta)$ .

As we've seen, if we wish to compute the thermal partition function  $Z = \text{Tr} e^{-\beta H}$  using the path integral then we must give the fermions anti-periodic boundary conditions. But, in doing so, we break supersymmetry. In contrast, if we wish to compute the Witten index  $\text{Tr} (-1)^F e^{-\beta H}$  then the path integral enjoys supersymmetry. This makes intuitive sense. In general, the full partition function  $Z$  is no easier to compute for a supersymmetric theory than a non-supersymmetric theory. But the Witten index is much easier. And, from the path integral perspective, this manifests itself because of the transformations (2.14).

To proceed, let's first show that the Witten index

$$\mathcal{I} = \text{Tr} (-1)^F e^{-\beta H} = \int \mathcal{D}x \mathcal{D}\psi^\dagger \mathcal{D}\psi e^{-S_E[x, \psi, \psi^\dagger]}$$

doesn't care about the magnitude of the potential. To this end, we rescale  $h \rightarrow \lambda h$  with  $\lambda > 0$ . We then differentiate with respect to  $\lambda$  to find

$$\frac{d\mathcal{I}}{d\lambda} = \int \mathcal{D}x \mathcal{D}\psi \mathcal{D}\psi^\dagger \left( - \oint d\tau (\lambda h'^2 - h'' \psi^\dagger \psi) \right) e^{-S_E}$$

The extra term in the integrand has a special form because it is itself a supersymmetry variation. To see this, it's useful to use the supersymmetry generators that we introduced in (1.21). With a rescaled potential  $\lambda h$  and Euclidean time, these become

$$\begin{aligned} \mathcal{Q}_\lambda &= \int dt \left[ \psi(t) \frac{\delta}{\delta x(t)} - \left( \frac{dx}{d\tau} - \lambda h' \right) \frac{\delta}{\delta \psi^\dagger(t)} \right] \\ \mathcal{Q}_\lambda^\dagger &= \int dt \left[ -\psi^\dagger(t) \frac{\delta}{\delta x(t)} + \left( \frac{dx}{d\tau} + \lambda h' \right) \frac{\delta}{\delta \psi(t)} \right] \end{aligned} \quad (2.15)$$

Then look at

$$\begin{aligned} \mathcal{Q}_\lambda^\dagger \oint d\tau h' \psi &= \oint d\tau \left( -h'' \psi^\dagger \psi + h' \frac{dx}{d\tau} + \lambda h'^2 \right) \\ &= \oint d\tau \left( -h'' \psi^\dagger \psi + \frac{dh}{d\tau} + \lambda h'^2 \right) \\ &= \oint d\tau \left( -h'' \psi^\dagger \psi + \lambda h'^2 \right) \end{aligned}$$

where, in the final term, we lost the total derivative. (Note that there's no danger of a boundary term here because  $\tau$  parameterises a circle and all fields are periodic.) This means that we can write the derivative of the Witten index as

$$\frac{d\mathcal{I}}{d\lambda} = \int \mathcal{D}x \mathcal{D}\psi \mathcal{D}\psi^\dagger \left( -\mathcal{Q}_\lambda^\dagger \oint d\tau h' \psi \right) e^{-S_E}$$

But we also know that the action is invariant under supersymmetry and, as we showed in (1.22), this can be written as  $\mathcal{Q}_\lambda^\dagger S_E = 0$ . This means that our final expression is a path integral of a total supersymmetry variation,

$$\frac{d\mathcal{I}}{d\lambda} = \int \mathcal{D}x \mathcal{D}\psi \mathcal{D}\psi^\dagger \mathcal{Q}_\lambda^\dagger \left( -e^{-S_E} \oint d\tau h' \psi \right)$$

The integrand is said to be *Q-exact*. The all-important point is that the integral of any Q-exact quantity always vanishes.

To see this, note from (2.15) that there are two terms in  $\mathcal{Q}^\dagger$  (or  $\mathcal{Q}$ ): one in which we differentiate with respect to  $x(t)$ , and one in which we differentiate with respect to  $\psi(t)$ . Let's start with the second of these.

To set the scene, we'll briefly return to normal Grassmann integration (as opposed to functional integration). Recall that the integral over any Grassmann variable  $\theta$  only gives a non-zero answer if there's a single copy of  $\theta$  in the integrand,

$$\int d\theta 1 = 0 \quad \text{and} \quad \int d\theta \theta = 1$$

There can't be more powers of  $\theta$  in the integrand because these are Grassmann variables and  $\theta^2 = 0$ . This means that Grassmann integration always obeys

$$\int d\theta \frac{d}{d\theta}(\text{Anything}) = 0 \tag{2.16}$$

That's because the  $d/d\theta$  kills any power of  $\theta$  that may have been lurking in the expression "Anything", ensuring that there's nothing left to saturate the  $\int d\theta$  integral. The formula (2.16) looks very much like an "integration by parts" formula for Grassmann variables, but with no danger of a boundary term.

The story above also holds for the functional integration over fermionic fields. We have

$$\int \mathcal{D}\psi \mathcal{D}\psi^\dagger \frac{\delta}{\delta\psi(t)}(\text{Anything}) = \int \mathcal{D}\psi \mathcal{D}\psi^\dagger \frac{\delta}{\delta\psi^\dagger(t)}(\text{Anything}) = 0$$

That deals with the fermionic functional derivatives in  $\mathcal{Q}$  and  $\mathcal{Q}^\dagger$ .

We're left with the bosonic functional derivatives  $\delta/\delta x(t)$ . Here we have a total derivative, albeit of a functional kind and we would expect such an integral to be given by the boundary term. The question is: what should we consider to be the boundary of this functional space? Large  $x(t)$ ? Wildly varying  $x(t)$ ? Either way, the boundary term vanishes. This is because there is an exponential suppression from the action  $e^{-S_E}$  that asymptotes quickly to zero for anything that you might reasonably consider to be the boundary. The upshot of these arguments is that

$$\int \mathcal{D}x \mathcal{D}\psi \mathcal{D}\psi^\dagger \mathcal{Q}^\dagger(\text{Anything}) = \int \mathcal{D}x \mathcal{D}\psi \mathcal{D}\psi^\dagger \mathcal{Q}(\text{Anything Else}) = 0$$

This, in turn, ensures that

$$\frac{d\mathcal{I}}{d\lambda} = 0$$

which, of course, we know to be true from our Hamiltonian analysis.

Now we're in business. Because the Witten index is independent of  $\lambda$ , we can calculate it in the limit  $\lambda \rightarrow \infty$ . Here the potential term in the action suppresses all contribution except for the a finite number of constant maps,

$$x(\tau) = X \quad \text{such that } h'(X) = 0$$

There are the critical points of  $h$ . The phenomenon of an integral – in this case an infinite dimensional functional integral – receiving contributions from just a handful of points is known as *localisation*. It is a property of supersymmetric path integrals that is not shared by most other quantum systems.

We still need to compute the partition function around each of these critical points. As we increase  $\lambda$ , the potential around each critical point gets steeper and steeper and the physics can be better and better approximated by a harmonic oscillator, with

$$h'(x) \approx h''(X)(x - X) + \dots \quad \Rightarrow \quad V(x) \approx \frac{1}{2}(h''(X))^2(x - X)^2 + \dots$$

Indeed, taking the  $\lambda \rightarrow \infty$  limit should be viewed as suppressing the non-linear interactions in the potential. The statement that the Witten index is independent of  $\lambda$  is equivalently to saying that the one-loop approximation is, in fact, exact.

All of this means that the path integral expression for the Witten index is

$$\text{Tr}(-1)^F e^{-\beta H} = \int \mathcal{D}x \mathcal{D}\psi \mathcal{D}\psi^\dagger e^{-S_E} = \sum_X \frac{\det(d/d\tau - h''(X))}{\det^{1/2}(-d^2/d\tau^2 + h''(X)^2)}$$

where the sum is over the critical points. Happily, we've already put some effort into computing the determinants of these operators. The bosonic contribution is (2.9)

$$\det^{1/2} \left( -\frac{d^2}{d\tau^2} + h''(X)^2 \right) = 2 \sinh \left( \frac{\beta |h''(X)|}{2} \right)$$

Meanwhile, the fermionic contribution is (2.12)

$$\det \left( \frac{d}{d\tau} - h''(X) \right) = -2 \sinh \left( \frac{\beta h''(X)}{2} \right)$$

The end result is

$$\text{Tr}(-1)^F e^{-\beta H} = \sum_X \frac{-h''(X)}{|h''(X)|} = \sum_X \text{sign}(-h''(X))$$

This is the answer we expected. If  $h(x)$  is a polynomial of odd degree, then it has an even number of critical points  $X$ , with  $h''(X)$  alternating in sign, giving  $\text{Tr}(-1)^F e^{-\beta H} = 0$ . Meanwhile, if  $h(x)$  is a polynomial of even degree then the alternating signs don't cancel out, leaving  $\text{Tr}(-1)^F e^{-\beta H} = \pm 1$ .

## 2.2 Instantons

Much of our story so far has revolved around understanding the structure of ground states in supersymmetric systems. A common theme – one familiar from other quantum mechanical models – is that the existence of multiple classical ground states does not necessarily mean that there are multiple quantum ground states.

In this section, we develop a more hands-on understanding of how ground states are lifted. Once again, our tool of choice will be the path integral and, as we will see, this provides a particularly direct way to think about quantum tunnelling and related phenomena. We will explore how this works in some detail, first in ordinary quantum mechanical systems and then in those that exhibit supersymmetry.

The path integral in Euclidean time is (2.3),

$$\langle x_f | e^{-HT} | x_i \rangle = \int_{x(-T/2)=x_i}^{x(T/2)=x_f} \mathcal{D}x(\tau) e^{-S_E[x(\tau)]} \quad (2.17)$$

To start, we'll focus only on the bosonic degrees of freedom and then introduce fermions into the discussion later. We'll also restrict attention to just a single degree of freedom  $x(\tau)$ , with Euclidean action

$$S_E = \int d\tau \left[ \frac{1}{2} \left( \frac{dx}{d\tau} \right)^2 + \frac{1}{2} \left( \frac{dh}{dx} \right)^2 \right] \quad (2.18)$$


Although the specific form of the potential  $V = \frac{1}{2}h'^2$  arises naturally in any supersymmetric theory, it is possible to write any positive definite potential in this way. Moreover, as we now show, this turns out to be a useful thing to do even in a non-supersymmetric theory.

Tunnelling is particularly easy to understand from the path integral perspective. It arises from paths that start at one minima and end up at another. If the parameters in the potential are such that we can do a semi-classical analysis, then the amplitude for tunnelling is dominated by the classical paths that minimise  $S_E$ . There is a rather cute way of finding these paths. We write the action (2.18) by completing the square

$$S_E = \int d\tau \frac{1}{2} \left( \frac{dx}{d\tau} \mp \frac{dh}{dx} \right)^2 \pm \frac{dx}{d\tau} \frac{dh}{dx}$$

The first term is positive definite, the second a total derivative. This means that we have

$$S_E \geq \pm \int d\tau \frac{dh}{d\tau} = \pm(h(x_f) - h(x_i)) \quad (2.19)$$



**Figure 5.** The instanton profile.

If we fix the end points  $x_i$  and  $x_f$  to be two distinct minima, then the action is minimised when this inequality is saturated with the most stringent  $\pm$  sign. This means that if  $h(x_f) > h(x_i)$ , we should solve the equation

$$\frac{dx}{d\tau} = \frac{dh}{dx} \quad (2.20)$$

Solutions to this equation are known as *instantons*. The name is chosen (by 't Hooft) to mimic the names give to particles but, as will see, these solutions are not localised in space but in (Euclidean) time and so occur just for an instant. If  $h(x_f) < h(x_i)$ , we should solve the other equation

$$\frac{dx}{d\tau} = -\frac{dh}{dx} \quad (2.21)$$

Solutions to this equations are called *anti-instantons*. They interpolate between the two vacua in the opposite direction to instantons.


It will be useful to look at an example. Suppose that we take

$$h = \frac{\omega}{2a} \left( \frac{1}{3}x^3 - a^2x \right) \quad \Rightarrow \quad V = \frac{\omega^2}{8a^2}(x^2 - a^2)^2 \quad (2.22)$$

This is a double well potential with minima at  $x = \pm a$ . The coefficient out front is chosen so that, around each minima, the potential is approximated by a harmonic oscillator with frequency  $\omega$ ,

$$V(x) \approx \frac{1}{2}\omega^2(x \pm a)^2 + \dots \quad (2.23)$$

We have  $h(-a) = \omega a^2/3 > h(a) = -\omega a^2/3$ . The instanton therefore interpolates from  $x = +a$  at  $\tau \rightarrow -\infty$  to  $x = -a$  to  $\tau \rightarrow +\infty$ . In this case, the solution to (2.20) is



**Figure 6.** On the left: a double well potential  $V(x)$  with two minima. On the right, the same potential but flipped to  $-V(x)$  as befits Euclidean time.

straightforward:

$$x_{\text{inst}}(\tau) = -a \tanh\left(\frac{\omega(\tau - \tau_1)}{2}\right) \quad (2.24)$$

The profile of the instanton is shown in Figure 5. As  $\tau \rightarrow \infty$ , we see that  $x(\tau) \approx \mp a e^{\mp\omega\tau} \rightarrow \mp a$  and the instanton asymptotes exponentially quickly back to the vacuum. The profile deviates significantly from the vacuum only in region of width  $\sim 1/\omega$ . The exact position  $\tau = \tau_1$  where this happens is an arbitrary integration constant.

For this example, the action of the instanton is

$$S_{\text{inst}} = \frac{2\omega a^2}{3}$$

For more general  $h(x)$  the exact solution of the instanton may be harder to come by but its simple to get an intuitive feel for its properties. Viewed from the usual perspective of Lagrangian dynamics, the Euclidean action (2.18) describes a particle moving in a potential  $-V(x)$ . This is shown on the right-hand side of Figure 6 for the double well potential. The instanton (or anti-instanton) describes a particle that starts at one maximum of  $-V(x)$  at  $\tau \rightarrow -\infty$  and then rolls down and up to another maximum, reaching the peak only at  $\tau \rightarrow +\infty$ .

If  $V(x)$  has multiple minima, then we can only find solutions to the instanton equations (2.20) and (2.21) that interpolate between *neighbouring* minima. This is because these are first order equations of motion, and once you sit at a critical point of  $h$  you have necessarily stopped. That doesn't mean that there is no tunnelling between multiple vacua: indeed, as we'll see shortly, in non-supersymmetric quantum mechanics it is usually approximate solutions to the classical equations of motion that dominate proceedings.

### 2.2.1 Tunnelling

To get a feel for the path integral manipulations, we won't yet introduce supersymmetry. Instead we'll explore the quantum mechanical physics of the double well potential shown in the left-hand side of Figure 6.

Let's first remind ourselves what we qualitatively expect from the ground states. Around each minima, the potential looks like a harmonic oscillator (2.23) and we can then construct approximations to the ground states as Gaussian wavefunctions, localised around each of the minima

$$\psi_{\text{left}}(x) = \exp\left(-\frac{\omega}{2}(x+a)^2\right) \quad \text{and} \quad \psi_{\text{right}}(x) = \exp\left(-\frac{\omega}{2}(x-a)^2\right)$$

For any even potential  $V(x) = V(-x)$ , the energy eigenstates are also eigenstates of the parity operator, meaning that they are either even or odd functions. A better approximation to the low-lying energy eigenstates must therefore be

$$\psi_{\pm}(x) \approx \psi_{\text{left}}(x) \pm \psi_{\text{right}}(x)$$

But the true ground state of any quantum system has no node, meaning that  $\psi(x) \neq 0$  for any finite  $x$ . (Given a wavefunction  $\psi(x)$  with a node, we can consider  $|\psi(x)|$  and then smooth out the cusp to lower the expected energy.) So it must be that  $\psi_+(x)$  is an approximation to the ground state, while  $\psi_-(x)$  is an approximation to the first excited state.

We'll now add some quantitative meat to these statements using the path integral which provides a particularly straightforward way to compute the ground state energies of the double well potential. To see this, we pick position eigenstates  $|x_i\rangle$  and  $|x_f\rangle$ . These need not themselves be ground states of the system, but should have a non-zero overlap with the ground states. As we've seen, the path integral (2.17) naturally computes  $\langle x_f | e^{-HT} | x_i \rangle$ . If we insert a complete set of energy eigenstates  $|n\rangle$ , with energy  $E_n$ , then we have

$$\langle x_f | e^{-HT} | x_i \rangle = \sum_n e^{-E_n T} \langle x_f | n \rangle \langle n | x_i \rangle$$

If we wait long enough, this sum is dominated by the ground state  $E_0 < E_n$  for all  $n \neq 0$ . We then have, for large  $T$ ,

$$\langle x_f | e^{-HT} | x_i \rangle \sim e^{-E_0 T}$$

So to determine the ground state energy, we just need to compute the path integral and extract the large  $T$  behaviour. We can then find the ground state energy in the exponent.

We will now use the semi-classical, or WKB, approximation to compute the amplitude for a particle to tunnel from one vacuum to the other and, in doing so, extract the ground state energy. We start by using the path integral to compute the amplitude for the particle to tunnel from one classical vacuum to another due to a single instanton. We write

$$x(\tau) = x_{\text{inst}}(\tau) + \delta x$$

where  $x_{\text{inst}}(\tau)$  is the solution to the relevant instanton equation (either (2.20) or (2.21)). We then expand the Euclidean action as

$$S_E[x] = S_{\text{inst}} + \int d\tau \frac{1}{2} \delta x \left( -\frac{d^2}{d\tau^2} + V'' \right) \delta x + \mathcal{O}(\delta x^3) \quad (2.25)$$

Here  $V''$  is evaluated on  $x_{\text{inst}}$ . Similarly,  $S_{\text{inst}}$  is the action of the instanton which, from (2.19), is

$$S_{\text{inst}} = |h(x_f) - h(x_i)| \quad (2.26)$$

Alternatively, written in terms of the potential  $V = \frac{1}{2}h'^2$ , the action of the instanton is  $S_{\text{inst}} = \int dx \sqrt{2V}$ . It should be thought of as a measure of the difficulty in getting up and over (or, more precisely, through) the barrier between the two minima.

The semi-classical approximation is valid whenever we can ignore the  $\mathcal{O}(\delta x^3)$  contributions relative to the  $\delta x^2$  contributions in the path integral. To understand the circumstances under which this holds, we should look more closely at the action and identify a dimensionless coupling constant  $g$  which multiplies all higher order terms. Perturbation theory is then valid when  $g \ll 1$ . A simpler way to view things is to rescale the potential  $h(x) \rightarrow \lambda h(x)$ . Then the semi-classical approximation is valid in the limit  $\lambda \gg 1$  where we have a steep potential, with deep minima. Under this rescaling, the action of the instanton (2.26) becomes

$$S_{\text{inst}} \rightarrow \lambda |h(x_f) - h(x_i)|$$

and so we see that  $\lambda \gg 1$  is equivalent to

$$S_{\text{inst}} \gg 1$$

This is the requirement that we will use for the semi-classical approximation to be valid. The results that we will get below will receive corrections of order  $1/S_{\text{inst}}$ .

In the language of quantum field theory, neglecting the higher order  $\delta x^3$  terms is tantamount to computing one-loop diagrams but not two-loop or higher. In normal circumstances, we would be doing perturbation theory around the classical vacuum  $x(\tau) = \pm a$ , in which case we would have  $V'' = \omega^2$ , a constant. The difference here is that we're now doing perturbation theory around the background of the instanton profile.

The kind of instanton calculations that we're performing here are often referred to as *non-perturbative*. This refers to the fact that tunnelling phenomena of this kind can't be captured by perturbation theory around any single vacuum. However, the phrase "non-perturbative" is also a little misleading: we're still doing perturbation theory, just around a non-trivial solution.

Inserting (2.25) into the path integral (2.17), and dropping the terms that are cubic or higher, we are left with a Gaussian integral

$$\langle -a|e^{-HT}|+a\rangle\Big|_{\text{one-inst}} = e^{-S_{\text{inst}}} \int \mathcal{D}\delta x \exp\left(-\int d\tau \frac{1}{2}\delta x \left(-\frac{d^2}{d\tau^2} + V''\right) \delta x + \dots\right)$$

On the left-hand side, we've taken the tunnelling to happen over a time  $T$ ; ultimately we will be interested in taking  $T \rightarrow \infty$ . We have also stressed that we're computing the contribution to the tunnelling from a single instanton and we'll subsequently see that this is just part of the story.

Now we're in a familiar situation. The Gaussian integral gives, as usual, by

$$\det^{-1/2}\left(-\frac{d^2}{d\tau^2} + V''\right) \tag{2.27}$$

As we stressed above, this differs from the usual determinant that we compute in perturbation theory only because  $V'' = V''(x_{\text{inst}}(\tau))$  is now evaluated on the time-dependent profile  $x_{\text{inst}}(\tau)$ . Nonetheless, the strategy to computing the determinant remains the same: we first find the eigenvalues

$$\left(-\frac{d^2}{d\tau^2} + V''\right) \delta x = \lambda \delta x \tag{2.28}$$

The determinant is then given by the (suitably regularised) product of eigenvalues  $\lambda$ .

There is, however, a catch. In the background of the instanton, there is always one eigenvalue that is zero. Viewed naively, this would seem to tell us that the determinant vanishes, giving an infinite amplitude for tunnelling. This, it turns out, is not an infinity that we should try to regulate away, but instead an infinity that means we should think more carefully about what we're calculating. Our first task, therefore, is to understand the physics behind this zero eigenvalue.

Happily, there is a simple reason for the existence of this zero eigenvalue. It follows from the fact that, as seen in the explicit instanton profile (2.24), the instantons come with an integration constant  $\tau_1$  which specifies the “time” at which the profile jumps from one ground state to the other. Clearly the action of the instanton  $x_{\text{inst}}(\tau - \tau_1)$  is independent of  $\tau_1$ . But this means that

$$\delta x_0 = \frac{\partial x_{\text{inst}}}{\partial \tau_1} \quad (2.29)$$

obeys (2.28) with a vanishing eigenvalue  $\lambda = 0$ .

Understanding zero modes is an important part of any instanton computation. They typically arise, as in the present case, because the instanton solution is not unique, but labelled by a number of parameters known as *collective coordinates*. For us, the instanton profile has a single collective coordinate,  $\tau_1$ . Any fluctuation, like (2.29), that can be thought of as a variation of a collective coordinate necessarily has zero eigenvalue. These fluctuations are called *zero modes*.

In the present case, the existence of the zero mode can be traced to the fact that the underlying quantum mechanics enjoys time translation symmetry, while any particular instanton profile does not. In quantum field theory (or statistical field theory), we would refer to the zero mode as a “Goldstone boson” for time translational symmetry.

Now that we understand that the zero mode simply corresponds to the possible times,  $-T/2 < \tau_1 < T/2$ , at which the instanton makes its move, it’s clearer how we should proceed. We should treat the zero mode separately. First we integrate over all the non-zero modes. Then, rather than attempting to integrate over the zero mode, we instead exchange this for an explicit integration over the collective coordinate  $\tau_1$ ; this will simply multiply our final expression by an overall factor of  $T$ , the time over which the tunnelling takes place. The end result is

$$\langle -a | e^{-HT} | +a \rangle \Big|_{\text{one-inst}} = e^{-S_{\text{inst}}} \int_{-T/2}^{T/2} \frac{d\tau_1}{\sqrt{2\pi}} J \sqrt{\frac{1}{\det'(-\partial_\tau^2 + V'')}} \quad (2.30)$$

There are a few things to unpick in this formula. First, this is only the one-loop contribution and, strictly speaking, we should include a  $+\dots$  corresponding to higher loop contributions. Next, the determinant is written as  $\det'$ , with the prime denoting that we include only non-vanishing eigenvalues. The  $\sqrt{2\pi}$  is the standard normalisation for each mode in the path integral.

Finally, there is that factor of  $J$ : this is merely the Jacobian that arises when changing from integrating over the field  $x(\tau)$ , to the collective coordinate  $\tau_1$ . It is easy to calculate

$$J^2 = \int d\tau \left( \frac{\partial x_{\text{inst}}}{\partial \tau_1} \right)^2 = \int d\tau \frac{1}{2} \left( \left( \frac{\partial x_{\text{inst}}}{\partial \tau} \right)^2 + \left( \frac{dh}{dx} \right)^2 \right) = S_{\text{inst}} \quad (2.31)$$

where, in the second equality, we've used the fact that  $x_{\text{inst}}(\tau)$  satisfies the instanton equation (2.20).

There is one further step that is useful. We will write our expression in way that makes the comparison to the classical ground state energy clearer. As we've seen, each classical ground state is given by a harmonic oscillator of frequency  $\omega$ . We already computed the path integral for Euclidean periodic time  $\beta$  in Section 2. The long time behaviour must be independent of the boundary conditions, so we also have

$$\langle a | e^{-HT} | a \rangle_{\text{SHO}} = \sqrt{\frac{1}{\det(-\partial_\tau^2 + \omega^2)}} \approx e^{-\omega T/2} \quad (2.32)$$


where  $e^{-\omega T/2}$  is the long time behaviour of the  $1/\sinh$  formula that we derived in (2.8). It will ultimately be clearer to write our tunnelling amplitude in a way that highlights the connection to the harmonic oscillator so, to this end, we collect everything together to get the final result for the one-instanton contribution,

$$\langle -a | e^{-HT} | +a \rangle \Big|_{\text{one-inst}} = T e^{-\omega T/2} K e^{-S_{\text{inst}}} \quad (2.33)$$

where all the other pre-factors have been bundled together into

$$K = \sqrt{\frac{S_{\text{inst}}}{2\pi}} \sqrt{\frac{\det(-\partial_\tau^2 + \omega^2)}{\det'(-\partial_\tau^2 + V'')}}$$

There are three things to take away from this. First, there are some slightly messy pre-factors that we've absorbed into  $K$ , which now include a ratio of the harmonic oscillator and instanton determinants. The exact expression for this ratio will not be particularly important in what follows and we won't make any attempt to compute it. However, the advantage of writing this as a ratio of determinants is that it makes it clear that it differs from 1 only due to physics in a region of width  $1/\omega$  where the instanton profile is non-trivial, and  $V''(x_{\text{inst}})$  differs from  $\omega^2$ . We'll see the utility of this shortly.



**Figure 7.** A dilute gas consisting of an instanton, followed by an anti-instanton and then, finally, another instanton.

Second, the amplitude is suppressed by a factor of  $e^{-S_{\text{inst}}}$ . This is a characteristic feature of tunnelling in quantum mechanics. Finally, we see that the tunnelling amplitude from a single instanton has the slightly odd  $Te^{-T}$  behaviour. It turns out that the correct interpretation of this comes by considering not a lone instanton, but a whole slew of them.

### 2.2.2 The Dilute Gas Approximation

In the calculation above, we restricted to a single instanton solution that interpolates from one classical ground state to the other. However, we know that the interesting part of this instanton profile takes place over a region that is exponentially localised within a width  $\sim 1/\omega$ . That means that if we take an instanton, followed a long time later, by an anti-instanton, followed some time later still by another instanton, then this *almost* solves the classical equation of motion. It's not an exact solution because there are no exact classical solution with these properties. But, if the instantons and anti-instantons are separated by a distance  $L \gg 1/\omega$ , then the action of a string of  $n$  such objects is roughly

$$S_{n\text{-inst}} = nS_{\text{inst}} + \mathcal{O}(e^{-\omega L})$$

This means that the action decreases very little as  $L$  increases. In this sense, as long as  $L \gg 1/\omega$ , the deviation from an exact solution is small.

Our interest in the classical instanton solutions is purely as a starting point for a semi-classical evaluation of the path integral. But, for these purposes, the approximate solutions, consisting of a string of instantons and anti-instantons are equally as good. This is known as the *dilute gas approximation*. An example of a dilute gas is shown in Figure 7.

We take the locations of these instantons and anti-instantons to lie at

$$-\frac{T}{2} < \tau_1 < \tau_2 < \dots < \tau_n < \frac{T}{2} \quad (2.34)$$

where  $\tau_k$  is the position of an instanton for  $k$  odd, and an anti-instanton for  $k$  even. The dilute gas approximation holds if  $\tau_{k+1} - \tau_k \gg 1/\omega$ .

In computing the amplitude  $\langle -a|e^{-HT}|a\rangle$ , we should sum over all possible numbers of instantons and anti-instantons. We just need one more instanton than anti-instanton to ensure that we end up in the opposite vacuum from where we started. In other words,  $n$  should be odd in (2.34).

Because the (anti)-instantons are far separated, their contribution to the path integral are independent. That means that we can simply import the calculation that we did above and the full tunnelling amplitude generalises the one-instanton result (2.33)

$$\langle -a|e^{-HT}|+a\rangle = e^{-\omega T/2} \sum_{n \text{ odd}} \int_{-T/2}^{T/2} d\tau_1 \int_{\tau_1}^{T/2} d\tau_2 \dots \int_{\tau_{n-1}}^{T/2} d\tau_n (K e^{-S_{\text{inst}}})^n$$

Note that the harmonic oscillator contribution  $e^{-\omega T/2}$  sits out the front of everything. Instead, each (anti)-instanton independently contributes a factor of the ratio of determinants  $K$  since, as we argued above, this ratio of determinants is non-trivial only in the vicinity of the (anti)-instanton.

The factor of  $T$  in (2.33), which came from the integral over the collective coordinate  $\tau_1$ , is now replaced by the multi-integral above. This is straightforward to evaluate and gives

$$\langle -a|e^{-HT}|+a\rangle = e^{-\omega T/2} \sum_{n \text{ odd}} \frac{T^n}{n!} (K e^{-S_{\text{inst}}})^n = e^{-\omega T/2} \sinh(KT e^{-S_{\text{inst}}})$$

We see the effect of summing over the dilute gas is to exponentiate the one-instanton contribution  $KT e^{-S_{\text{inst}}}$ .

We can also do a similar calculation to evaluate the amplitude  $\langle +a|e^{-HT}|+a\rangle = \langle -a|e^{-HT}| - a\rangle$  for returning to our original vacuum. Everything is the same, except that we should now take the number  $n$  of instantons and anti-instantons to be even. Of course,  $n = 0$  is allowed. We then get

$$\langle -a|e^{-HT}| - a\rangle = e^{-\omega T/2} \sum_{n \text{ even}} \frac{T^n}{n!} (K e^{-S_{\text{inst}}})^n = e^{-\omega T/2} \cosh(KT e^{-S_{\text{inst}}})$$

Before we go on, we note that this same calculation appears in a seemingly different setting of [Statistical Field Theory](#) when we showed that discrete symmetries in 1d hot systems cannot be spontaneously broken. (See section 1.3.3 of those notes.)

The two formulae above contain the information about the energy splitting that we wanted to find. From our earlier discussion, we know that the ground state has non-vanishing overlap with

$$|\text{ground}\rangle = | + a\rangle + | - a\rangle$$

while the first excited state has overlap with

$$|\text{excited}\rangle = | + a\rangle - | - a\rangle$$

From above, we have

$$\langle \text{ground} | e^{-HT} | \text{ground} \rangle = 2e^{-E_0 T} \quad \text{with} \quad E_0 = \frac{\omega}{2} - Ke^{-S_{\text{inst}}}$$

and

$$\langle \text{excited} | e^{-HT} | \text{excited} \rangle = 2e^{-E_1 T} \quad \text{with} \quad E_1 = \frac{\omega}{2} + Ke^{-S_{\text{inst}}}$$

We see the promised energy splitting, proportional to the characteristic tunnelling amplitude  $e^{-S_{\text{inst}}}$ .

Strictly speaking, neither of the formulae above can be trusted. Both  $E_0$  and  $E_1$  will receive perturbative contributions to their energies and these will scale as some power of  $1/S_{\text{inst}}$ . The important fact is that, because of the symmetry of the potential, these contributions will be the same for both states. The real meaning of the calculation we've just done is to compute the splitting of the two states

$$E_1 - E_0 = 2Ke^{-S_{\text{inst}}}$$

Of course, if we really want to do a good job then we should roll up our sleeves and compute the ratio of determinants that sits in  $K$ . But we can see the key piece of physics without doing this: the splitting of energy levels scales as  $e^{-S_{\text{inst}}}$ .

### 2.3 Instantons and Supersymmetry

It's now time to return to supersymmetric quantum mechanics. It turns out that there is a deep relationship between instantons and supersymmetry, both in quantum mechanics and in higher dimensional quantum field theories. The two make for perfect bedfellows. In this section, we will start to get a hint of where this relationship emerges from. We'll also see that the existence of fermions brings some important technical differences to the tunnelling calculation that we did in the last section.

For concreteness, we'll again work with the cubic  $h$  given in (2.22), corresponding to a double well potential  $V(x)$  with minima at  $x = \pm a$ . The novelty, of course, is the presence of fermions.

The Euclidean supersymmetric action is (2.13)

$$S_E[x, \psi, \psi^\dagger] = \int d\tau \left[ \frac{1}{2} \left( \frac{dx}{d\tau} \right)^2 + \psi^\dagger \frac{d\psi}{d\tau} + \frac{1}{2} h'^2 - h'' \psi^\dagger \psi \right] \quad (2.35)$$

We can largely proceed as the previous section. The bosonic instanton configuration is  $x_{\text{inst}}(\tau)$  and we evaluate the path integral in a semi-classical expansion around this background. In addition to the bosonic fluctuations  $\delta x$ , we must also integrate out the fermions. This give the usual determinant contribution

$$\det \left( \frac{d}{d\tau} - h'' \right) \quad (2.36)$$

where, as for the bosonic fluctuations, we evaluate  $h'' = h''(x_{\text{inst}}(\tau))$  on the instanton profile.

We need to briefly pause to think about what this determinant means because, in contrast to the bosonic fluctuations (2.27), it's not the determinant of a Hermitian operator. The operator used to be Hermitian, back when we were living in real time, where it was  $(+id/dt - h'')$ . But the Wick rotation ruined that property. We define

$$D = \frac{d}{d\tau} - h'' \quad \text{and} \quad D^\dagger = -\frac{d}{d\tau} - h'' \quad (2.37)$$

The eigenvalue equation of this pair of operators should be thought of as

$$Df(\tau) = \lambda g(\tau) \quad \text{and} \quad D^\dagger g(\tau) = \lambda f(\tau)$$

The determinant (2.36) is then the product of all eigenvalues  $\lambda$ .

### 2.3.1 Fermi Zero Modes

When we come to evaluate the fermionic determinant we run into the same subtlety that we saw in the bosonic case: the operator  $D$  has a zero eigenvalue and so the determinant is zero. In fact, the profile of the associated fermionic fluctuation takes the same form as the bosonic zero mode (2.29),

$$\psi_0 = \eta \frac{dx_{\text{inst}}}{d\tau} \quad (2.38)$$

Here  $\eta$  is a constant Grassmann parameter. You should think of it as a Grassmann collective coordinate, analogous to the bosonic collective coordinate  $\tau_1$ . It is straightforward to see that  $\psi_0$  is an eigenfunction with vanishing eigenvalue. We have

$$\frac{dx_{\text{inst}}}{d\tau} = \frac{dh}{dx} \quad \Rightarrow \quad \frac{d^2x_{\text{inst}}}{d\tau^2} - h'' \frac{dx_{\text{inst}}}{d\tau} = 0 \quad \Rightarrow \quad \left( \frac{d}{d\tau} - h'' \right) \psi_0 = 0$$

We could have anticipated the existence of this fermionic zero mode on symmetry grounds. Recall that we could trace the bosonic collective coordinate  $\tau_1$  to time translational symmetry since, while the action is invariant under time translations, any given instanton profile is not. Similarly, the fermionic collective coordinate can be traced to a fermionic symmetry which is, of course, supersymmetry. If we look again at the transformation rules (2.14) for the fermions in Euclidean time, then we see something rather nice:

$$\delta\psi = \epsilon \left( \frac{dx}{d\tau} + h' \right) \quad , \quad \delta\psi^\dagger = \epsilon^\dagger \left( -\frac{dx}{d\tau} + h' \right) \quad (2.39)$$

The supersymmetry transformations have, hidden within them, the instanton and anti-instanton equations (2.20) and (2.21)! This, it turns out, is a beautiful feature of supersymmetry, and one that persists as we look both to more complicated theories and to more complicated instantons and other solitons. For now, we note that if we take an instanton obeying  $\dot{x} = h'$  and hit it with a supersymmetry transformation, then  $\psi$  will turn on while  $\psi^\dagger$  will not. But, because supersymmetry is a symmetry, the action of the solution doesn't change when  $\psi$  turns on. This is the fermi zero mode (2.38) that we identified above.

You might be nervous that we seem to have broken reality. In the background of an instanton, the fermion  $\psi$  has a zero mode, but  $\psi^\dagger$  does not. Indeed, the equation of motion for  $\psi^\dagger$  is  $D^\dagger\psi^\dagger = 0$  and  $D^\dagger$  has no zero mode in the background of an instanton. Conversely, in the background of an anti-instanton  $\psi^\dagger$  has a zero mode, while  $\psi$  has none. This issue is commonplace for fermions in Euclidean time (or, more generally, in Euclidean space) and arises because  $D$  is not Hermitian. It's best to think of  $\psi$  and  $\psi^\dagger$  as independent degrees of freedom in Euclidean time. Only when we Wick rotate back to real time (or Minkowski space) do the reality and Hermiticity properties of various operators manifest themselves again.

The upshot is that the instanton breaks one half of the supersymmetries:  $Q^\dagger$  is broken and generates a fermionic zero mode, while  $Q$  survives. Objects, like instantons, which have the property of preserving some fraction of the supersymmetry are known as *BPS*. (The initials stand for Bogomolnyi, Prasad and Sommerfeld, but what they actually

did was only vaguely related to supersymmetry and the meaning of the initials BPS has evolved over the years.)

Although  $Q$  is unbroken, it is not totally redundant. It actually relates the collective coordinates  $\tau_1$  and  $\eta$  of the instanton, a kind of “zero dimensional” supersymmetry. This interpretation won’t be important for now.

Let’s now return to our tunnelling computation. We know what to do: rather than integrate over all fermion modes, we isolate the zero mode and treat it separately, choosing instead to integrate over the fermionic collective coordinate  $\eta$ . As before, we pick up a Jacobian factor which, because the fermion zero mode (2.38) has the same functional form as the bosonic zero mode (2.29), is the same value  $J = \sqrt{S_{\text{inst}}}$  that we computed in (2.31). But Jacobians for Grassmann integration come as  $1/J$ , rather than  $J$  so this actually cancels our original bosonic contribution.

The net effect is that if we repeat the steps that took us to (2.30), we now have

$$\langle -a|e^{-HT}|+a\rangle\Big|_{\text{one-inst}} = e^{-S_{\text{inst}}} \int_{-T/2}^{T/2} \frac{d\tau_1}{\sqrt{2\pi}} \int d\eta \frac{\det'(\partial_\tau - h'')}{\sqrt{\det'(-\partial_\tau^2 + V'')}}$$

We now have a ratio of determinants, both with zero eigenvalues omitted. There is no need to do our previous trick of introducing the harmonic oscillator amplitude (2.32). Indeed, part of the reason for doing that previously was to make manifest the  $\frac{1}{2}\hbar\omega$  ground state energy but, as we’ve seen, the analogous semi-classical energy in supersymmetric quantum mechanics is exactly zero.

### 2.3.2 Computing Determinants

In non-supersymmetric theories, it can be very challenging to compute the determinants in the background of an instanton. In contrast, in supersymmetric theories it is trivial because the ratio of determinants precisely cancels! To see this, we use the definition of the fermionic operators in (2.37) and note that

$$\begin{aligned} D^\dagger D &= \left(-\frac{d}{d\tau} - h''\right) \left(\frac{d}{d\tau} - h''\right) \\ &= -\frac{d^2}{d\tau^2} + h''' \frac{dx}{d\tau} + (h'')^2 \\ &= -\frac{d^2}{d\tau^2} + h''' h' + (h'')^2 \end{aligned}$$

where, to get to the second line, we’ve used the fact that these operators are evaluated on the solution to the instanton equation (2.20). But the potential is  $V = \frac{1}{2}h'^2$ , so

$V' = h'h''$  and  $V'' = h'''h' + (h'')^2$ , so

$$D^\dagger D = -\frac{d^2}{d\tau^2} + V''$$

which is precisely the bosonic fluctuation operator. This means that if we have a bosonic eigenfunction  $f$ , with  $D^\dagger Df = \lambda f$  with  $\lambda > 0$ , then we can define  $g = Df/\sqrt{\lambda}$ . We then have  $Df = \sqrt{\lambda}g$  and  $D^\dagger g = \sqrt{\lambda}f$ , which means that  $\sqrt{\lambda}$  is an eigenvalue of the fermionic operator (in the sense that we described previously).

This cancellation is entirely analogous to our previous observation that the ground state energy in a supersymmetric vacuum is zero, since the  $+\frac{1}{2}\hbar\omega$  from the harmonic oscillator is precisely cancelled by a  $-\frac{1}{2}\hbar\omega$  from the fermions. Here we see a similar cancellation persists about a BPS instanton configuration. This is a lesson that also transfers to higher dimensional quantum field theories, where it is often the case that all perturbative contributions cancel between bosons and fermions when evaluated about BPS backgrounds.

In the present context, it means that the tunnelling amplitude in a supersymmetric theory due to a single instanton is extremely simple:

$$\langle -a|e^{-HT}|+a\rangle\Big|_{\text{one-instanton}} = e^{-S_{\text{inst}}} \int_{-T/2}^{T/2} \frac{d\tau_1}{\sqrt{2\pi}} \int d\eta$$

Not only is it very simple, it is also very zero. That's because of the presence of the fermion zero mode. Recall the rules for Grassmann integration,

$$\int d\eta \ 1 = 0 \quad \text{and} \quad \int d\eta \ \eta = 1$$

With nothing to soak up the fermion zero mode in the integrand, the amplitude for tunnelling vanishes.

In fact, this is to be expected given our earlier discussion of supersymmetric quantum mechanics. From Section 1.2, we know that the semi-classical ground states  $|-a\rangle$  and  $|+a\rangle$  lie in different spin sectors or, equivalently, in different components of the Hilbert space factorisation  $\mathcal{H} = \mathcal{H}_B \oplus \mathcal{H}_F$ . This means that there can be no tunnelling from one state to another and the path integral realises this by introducing a lone fermion zero mode.

### 2.3.3 Computing the Ground State Energy

The Hamiltonian analysis of Section 1.2 told us more about this system. We know, for example, that the two states localised in different minima remain true ground states of the system but their energy is lifted above zero (i.e. supersymmetry is broken for a cubic  $h(x)$ ). It is possible to see this from the path integral. We just need a small tweak of our previous analysis.

Rather than working with position eigenstates  $|\pm a\rangle$ , we'll instead revert briefly to the exact ground states  $|L\rangle$  and  $|R\rangle$ , which have support localised around the left and right minima respectively. Supersymmetry means that these must have the same energy  $E_0$  and sit in  $\mathcal{H}_B$  and  $\mathcal{H}_F$  respectively. This means that  $Q^\dagger|R\rangle = Q|L\rangle = 0$ . Moreover, from our early analysis (1.3) we know that the two states are related by  $|R\rangle = Q^\dagger|L\rangle/\sqrt{2E_0}$ . The energy of either state can then be computed as follows:

$$E_0 = \langle L|H|L\rangle = \frac{1}{2}\langle L|\{Q, Q^\dagger\}|L\rangle = \frac{1}{2}\langle L|QQ^\dagger|L\rangle = \sqrt{\frac{E_0}{2}}\langle L|Q|R\rangle$$

This means that

$$E_0 = \frac{1}{2}|\langle L|Q|R\rangle|^2 \tag{2.40}$$

We see that, to compute the energy of the ground state, we must compute a tunnelling amplitude  $\langle L|Q|R\rangle$  but, crucially, with a factor of the supercharge  $Q$  sandwiched between the two states.

In fact, it turns out that there's a little trick and things work out better if we compute the amplitude  $\langle L|[Q, h]|R\rangle$ <sup>1</sup>. This is very closely related to the amplitude  $\langle L|Q|R\rangle$  that we need. First, for a steep potential, we have  $|R\rangle \approx | + a\rangle$  and  $|L\rangle \approx | - a\rangle$ , so  $h(x)|R\rangle \approx h(a)|R\rangle$  and  $h(x)|L\rangle = h(-a)|L\rangle$  and

$$\langle L|[Q, h]|R\rangle \approx (h(a) - h(-a))\langle L|Q|R\rangle = S_{\text{inst}} \langle L|Q|R\rangle$$

But the commutator  $[Q, h]$  has a particularly nice form. After Wick rotating the supercharge (1.17) reads  $Q = i(p - h')\psi$  and the commutator is

$$[Q, h] = \frac{dh}{dx} \psi$$

---

<sup>1</sup>This sidesteps an annoying subtlety. If you compute the matrix element for  $\langle L|Q|R\rangle$  directly then, at leading order, the result will vanish. This is because, after Wick rotation to Euclidean time,  $Q$  is proportional to the instanton equations and so vanishes when evaluated on the instanton. (This follows from the fact that the supersymmetry transformation (2.39) is proportional to the instanton equation.) You then have to work to higher order to find the non-vanishing ground state energy. Computing the matrix element of  $[Q, h]$  avoids this headache.

This means that we can get to the ground state energy (2.40) by computing the amplitude

$$\langle L|Q|R\rangle \approx \frac{1}{S_{\text{inst}}} \langle L|[Q, h]|R\rangle = \frac{1}{S_{\text{inst}}} \langle L|h'\psi|R\rangle$$

Now we can revert to our path integral expression again. We compute  $\langle L|Q|R\rangle$  by the same kind of analysis that we performed above, but this time with an extra power of  $h'\psi$  in the integrand,

$$\langle L|Q|R\rangle = \frac{e^{-S_{\text{inst}}}}{S_{\text{inst}}} \int_{-T/2}^{T/2} \frac{d\tau}{\sqrt{2\pi}} \int d\eta h' \psi$$

We next replace the  $\psi$  that appears in this expression with the fermi zero mode (2.38)  $\psi_0 = \eta dx/d\tau$ . Furthermore,  $h'$  should be evaluated on the the instanton background  $x_{\text{inst}}(\tau)$ . Importantly, the presence of  $\psi_0$  soaks up the  $\int d\eta$  integral, rescuing the result from the vanishing answer we found before. We now have

$$\langle L|Q|R\rangle = \frac{e^{-S_{\text{inst}}}}{S_{\text{inst}}} \int \frac{d\tau}{\sqrt{2\pi}} \frac{dh}{dx} \frac{dx}{d\tau} = \frac{e^{-S_{\text{inst}}}}{S_{\text{inst}}} \int \frac{d\tau}{\sqrt{2\pi}} \frac{dh}{d\tau}$$

Rather wonderfully, the final integral is a total derivative and just gives us  $h(\tau = +\infty) - h(\tau = -\infty) = S_{\text{inst}}$ . The final answer is then very simple:

$$\langle L|Q|R\rangle = \frac{e^{-S_{\text{inst}}}}{\sqrt{2\pi}} \tag{2.41}$$

We learn that the ground state energy is non-zero, but exponentially small

$$E_0 \sim e^{-2S_{\text{inst}}}$$

There's another lesson lurking in the calculation above. To compute the energy  $E_0$ , we didn't need to invoke the dilute gas approximation; it was sufficient to look at a single instanton. Indeed, viewed the right way it was *necessary* to look at just a single instanton. This is because the single instanton is BPS, meaning that it is invariant under one-half of the supersymmetries, and therefore has just a single fermion zero mode. However, a string of instanton-anti-instanton pairs does not have this property: it breaks both  $Q$  and  $Q^\dagger$  and therefore has two fermion zero modes, rather than just one. This is a special property of BPS instantons in supersymmetric theories that is closely related to the localisation of the path integral that we saw previously.

We'll revisit instanton calculations of this kind in Section 3.2 where we discuss Morse theory. It will turn out that these kind of calculations underlie many of the key ideas in that context.

### 2.3.4 One Last Example: A Particle on a Circle

Before we move on to more geometrical things, there is one last example that will prove useful to have under our belts. This is the supersymmetric particle moving on a circle  $S^1$ , with

$$h(x) = \omega R \sin(x/R)$$

where  $R$  is the radius of the circle. The associated potential is

$$V(x) = \frac{\omega^2}{2} \cos^2(x/R)$$

and has two minima at  $x = \pm\pi R/2$ .

We briefly discussed this model in Section 1.2.2 where we showed that, despite its similarities to the double well potential, it actually has two zero energy ground states, given by  $e^{+h}|0\rangle$  and  $e^{-h}\psi^\dagger|0\rangle$ . The puzzle that we'd like to address here is: why aren't these states lifted from the perspective of the path integral?

It's straightforward to guess the reason for this, but a little trickier to show how it works. Consider instantons (as opposed to anti-instantons) that solve

$$\frac{dx}{d\tau} = h' = \omega \cos(x/R)$$

These necessarily interpolate from small  $h(x)$  to large  $h(x)$  which, for us, means from the vacuum  $x = -\pi R/2$  at  $\tau \rightarrow -\infty$  to the vacuum  $x = +\pi R/2$  at  $\tau \rightarrow +\infty$ . The novelty is that we have two different instanton solutions in this case, corresponding to the two different ways to go around the circle. The first instanton has  $\dot{x} > 0$ , the second  $\dot{x} < 0$ .

So it's clear what the solution to our puzzle must be. These two instantons must contribute with opposite signs, so that they cancel out in the matrix element

$$\langle +\frac{\pi R}{2} | Q | -\frac{\pi R}{2} \rangle$$

that we care about, leaving the energy of both states at zero. The question is: how does this minus sign arise in the computation?

This, it turns out is subtle. A rerun of the calculation above shows that there's nowhere obvious that this sign could appear. The non-obvious place is, it turns out, in the definition of the determinants. The cancellation that we derived in Section 2.3.2 is really

$$\frac{\det D}{\sqrt{\det D^\dagger D}} = \pm 1$$

Figuring out which sign we get is not so straightforward. For now, we'll content ourselves with the observation that, by answer analysis, the signs must be opposite for the two instantons that traverse the circle in different directions. We'll give a prescription for computing this sign in Section 3.2 when we discuss Morse theory.

### 3 Supersymmetry and Geometry

In this section, we will begin our journey into the territory of mathematicians. Our strategy is to think about the physics of a particle moving on a manifold. As this section progresses, we will learn that the quantum ground states of this particle encode some precious information about the manifold.

Before we get to supersymmetry, let's set the scene. We consider a massive, non-relativistic particle moving on the manifold  $M$  of dimension  $\dim(M) = n$ . The dynamics of this particle is described by the Lagrangian

$$L = \frac{1}{2} g_{ij}(x) \dot{x}^i \dot{x}^j \quad (3.1)$$

where  $x^i$  are coordinates on the manifold, with  $i = 1, \dots, n$ , and  $g_{ij}(x)$  is a Riemannian metric on  $M$ .

Lagrangians of the form (3.1) are commonplace in physics, both in quantum mechanics and in higher dimensional quantum field theories. They often go by the unhelpful name of a *sigma model*. Sometimes they are called non-linear sigma models to reflect the fact that, unless  $g_{ij}$  is constant, the equations of motion will be non-linear. The name “sigma model” is utterly unilluminating; it dates from one of the first such models written down by Gell-Mann and Levy to describe the dynamics of mesons. (Somewhat comically, Gell-Mann and Levy were building on an earlier model that described both pions and an extra meson known as the “sigma”. They then wrote down an improved model that described just the mesons but chose to name it after the missing particle. And the name stuck.)

Geometrically, we should think of the sigma model as a map from the worldline of the particle  $W$  to the manifold,

$$x(t) : W \mapsto M$$

The manifold  $M$  is known as the *target space*. For much of what we do below, the story will be simplest if  $M$  is a compact, orientable manifold and we'll assume this to be the case in what follows.

Strictly speaking, the metric  $g_{ij}(x)$  in the Lagrangian should be viewed as the pull back of the metric from  $M$  to  $W$ . As we saw in [earlier courses](#) covering differential geometry, strictly speaking the sigma model only describes the particle in a patch of the manifold  $M$  that is covered by the coordinates  $x^i$ . One might think that to understand more subtle topological issues, we should be willing to consider overlapping patches. Perhaps surprisingly, it will turn out that this is not necessary, at least in these lectures.

We now ask: what does the particle described by (3.1) know about the manifold  $M$ , and what kind of mathematics might it encapsulate? To get a sense for this, we could first think about the Lagrangian (3.1) as describing a classical particle. In this case the equations of motion are the geodesic equations

$$\ddot{x}^i + \Gamma_{jk}^i \dot{x}^j \dot{x}^k = 0 \quad (3.2)$$

where  $\Gamma_{jk}^i$  is the Levi-Civita connection,

$$\Gamma_{jk}^i = \frac{1}{2} g^{il} (\partial_j g_{kl} + \partial_k g_{jl} - \partial_l g_{jk})$$

and we're using the notation  $\partial_i = \partial/\partial x^i$ .

There is certainly a lot of interesting physics in the geodesic equation. But it's challenging to extract any interesting mathematical statements about the manifold  $M$  from knowledge of these geodesics. In particular, at any given time, the particle knows only about its immediate surrounding, yet any point looks much the same as any other locally. This means that the state of the particle cannot know anything about the global properties of the manifold. To extract any such information, we would need to know about the entire history of the particle.

This can be contrasted with the situation in quantum mechanics. Now the wavefunction spreads over the manifold  $M$ , which suggests that the state of the particle may well know about some of the manifold's quirks. In particular, the state of a quantum particle may be sensitive to the topology of  $M$ . Ultimately, we will see that this is indeed the case, at least when we consider supersymmetric extension of our theory. But, for now, let's push on and consider the quantum theory associated to the non-supersymmetric Lagrangian (3.1).

To describe the quantum theory, we first need the momentum

$$p_i = \frac{\partial L}{\partial \dot{x}^i} = g_{ij} \dot{x}^j$$

We then impose canonical commutation relations  $[x^i, p_j] = i\delta_j^i$  and construct the Hamiltonian

$$H = p_i \dot{x}^i - L = \frac{1}{2} g^{ij} p_i p_j$$

Already here, things are not so straightforward because the metric  $g_{ij}$  depends on  $x^i$  and these don't commute with  $p_i$ . Different choices of ordering give different quantum Hamiltonians and so different theories.

There is no right or wrong choice here. But we can narrow down our options by requiring that the resulting theory has certain desirable properties. Given that we're interested in the geometry of  $M$ , it makes sense to search for a Hamiltonian that is covariant with respect to changes of coordinates on  $M$ . In other words, to stick as closely as possible to differential geometry. The action of the momentum on the wavefunction is, as usual,  $p_i = -i\partial_i$ , so the Hamiltonian should be a second order differential operator with terms that involve no more than two derivatives acting on the metric. There is a one-parameter family of such Hamiltonians, labelled by  $\alpha \in \mathbb{R}$ ,

$$H = -\frac{1}{2\sqrt{g}} \frac{\partial}{\partial x^i} \left( \sqrt{g} g^{ij} \frac{\partial}{\partial x^j} \right) + \alpha R \quad (3.3)$$

where  $g = \det g_{ij}$  and  $R$  is the Ricci scalar. The first term in this expression is the Laplacian, acting on functions, and can also be written more simply using the covariant derivative,

$$H = -\frac{1}{2} g^{ij} \nabla_i \nabla_j + \alpha R \quad (3.4)$$

We should also decide what Hilbert space we want our operators to act on. The obvious choice is to take the wavefunctions  $\psi(x)$  as functions over  $M$ , with the norm given by

$$\|\psi\|^2 = \int d^n x \sqrt{g} |\psi(x)|^2 \quad (3.5)$$

Note, in particular, that the inner product includes the factor of  $\sqrt{g}$  in the measure, as is appropriate in the geometric context.

Now we have our Hamiltonian (3.4) describing a quantum particle roaming around on a manifold  $M$ . What do we do with it? As physicists, our natural inclination is to find the spectrum of the Hamiltonian. We would typically expect that the particle has a unique ground state, with an infinite tower of excited states. This prompts two interesting questions: first, is it possible to calculate this spectrum? Second, what can we do with this information?

Both of these questions are interesting, although neither is easy. In general, it is a difficult problem to determine the spectrum of the Hamiltonian (3.4). Which properties of the manifold can be reconstructed from this spectrum is reminiscent of the famous question “can you hear the shape of a drum?”. Mathematicians have spent much time on this question. It is known, for example, that two manifolds may have the same spectrum even though they are not isometric. The first examples are 16-dimensional tori, but subsequent examples have been found in any dimension  $n \geq 2$ . In fact, it's

known that two manifolds may share the same spectrum even if they have different topology (e.g. their fundamental group may be different). All of which is to say that the problem of a quantum particle moving on a manifold  $M$  is certainly interesting, but thinking as a physicist provides no particular advantage. We will now see that this situation changes (for the better!) when we introduce supersymmetry.

### 3.1 The Supersymmetric Sigma Model

There is a beautiful generalisation of the sigma model Lagrangian (3.1) that admits supersymmetry. In addition to the  $n$  coordinates  $x^i$ , we also introduce  $n$  complex Grassmann variables  $\psi^i$ , and then consider the action

$$S = \int dt \frac{1}{2} g_{ij}(x) \dot{x}^i \dot{x}^j + i g_{ij} \psi^{\dagger i} \nabla_t \psi^j - \frac{1}{4} R_{ijkl} \psi^i \psi^j \psi^{\dagger k} \psi^{\dagger l} \quad (3.6)$$

The index  $i$  on  $\psi^i$  is telling us that the fermions live in the tangent space (strictly the tangent bundle) of  $M$ . This is highlighted by the appearance of the covariant derivative, pulled back to the worldline, in the fermion kinetic term

$$\nabla_t \psi^i = \frac{d\psi^i}{dt} + \Gamma_{jk}^i \frac{dx^j}{dt} \psi^k$$

As the particle moves on  $M$ , the fermions rotate due to this extra term. Finally, note that the four fermion term contracts with the Riemann tensor  $R_{ijkl}$ . This is the first suggestion that there might be some pretty geometry lurking in this theory. It's sometimes useful to note that the last term can also be written as

$$\frac{1}{4} R_{ijkl} \psi^i \psi^j \psi^{\dagger k} \psi^{\dagger l} = -\frac{1}{2} R_{ijkl} \psi^i \psi^{\dagger j} \psi^k \psi^{\dagger l}$$

The equivalence of these two expressions follows from the Riemann tensor identity  $R_{i[jkl]} = 0$ .

The action (3.6) is invariant under  $N = 2$  supersymmetry, given by the following supersymmetry transformations,

$$\begin{aligned} \delta x^i &= \epsilon^\dagger \psi^i - \epsilon \psi^{\dagger i} \\ \delta \psi^i &= \epsilon (-i \dot{x}^i + \Gamma_{jk}^i \psi^{\dagger j} \psi^k) \\ \delta \psi^{\dagger i} &= \epsilon^\dagger (+i \dot{x}^i + \Gamma_{jk}^i \psi^{\dagger j} \psi^k) \end{aligned} \quad (3.7)$$

The associated supercharges are:

$$Q = g_{ij} \dot{x}^i \psi^{\dagger j} \quad \text{and} \quad Q^\dagger = g_{ij} \dot{x}^i \psi^j \quad (3.8)$$

Note that, in contrast to Section 1, we have taken the supercharge  $Q$  to depend on  $\psi^\dagger$  rather than  $\psi$ . This is a notational convenience whose advantage we will see as we go along.

## How to Show that the Sigma Model is Supersymmetric

Conceptually, it's straightforward to demonstrate the supersymmetrictness of the sigma model: you just vary the action, use the transformations (3.7), and show that it vanishes. In practice, you end up with a tsunami of terms. Here's some help to guide you along the way.

First, when implementing the supersymmetry transformation it's useful to set  $\epsilon = 0$  and just keep the  $\epsilon^\dagger$  terms in the variation. There's no subtlety here: it's just means that we only have to keep track of half the terms in the variation. The other half are then fixed by ultimately requiring that the action and its variation are real. In particular, setting  $\epsilon = 0$  means that we have  $\delta\psi = 0$  while  $\delta\psi^\dagger \neq 0$ .

Second, there's a familiar trick, described in the lectures on [Quantum Field Theory](#), that is used to compute the conserved charges associated to any symmetry: we do local variations, instead of global variations. To this end we promote  $\epsilon^\dagger \rightarrow \epsilon^\dagger(t)$ . We will then find the supercharges multiplying the  $\epsilon^\dagger$  terms in the variation of the action.

Now we can start. Varying the action with  $\delta\psi = 0$  but  $\delta x, \delta\psi^\dagger \neq 0$  gives

$$\begin{aligned} \delta S = \int dt \, g_{ij} & \left( \dot{x}^i \delta \dot{x}^j + i \delta \psi^{\dagger i} \dot{\psi}^j + i \delta \psi^{\dagger i} \Gamma_{kl}^j \dot{x}^k \psi^l + i \psi^{\dagger i} \delta \Gamma_{kl}^j \dot{x}^k \psi^l + i \psi^{\dagger i} \Gamma_{kl}^j \delta \dot{x}^k \psi^l \right) \\ & + \delta g_{ij} \left( \frac{1}{2} \dot{x}^i \dot{x}^j + i \psi^{\dagger i} \dot{\psi}^j + i \Gamma_{kl}^j \dot{x}^k \psi^{\dagger i} \psi^l \right) \\ & - \frac{1}{4} \delta R_{ijkl} \psi^i \psi^j \psi^{\dagger k} \psi^{\dagger l} - \frac{1}{2} R_{ijkl} \psi^i \psi^j \delta \psi^{\dagger k} \psi^{\dagger l} \end{aligned}$$

where we've used  $R_{ijkl} = R_{ij[kl]}$  in the final term. Next it's useful to tame the terms by counting the number of fermions that they contain. There will be terms with 1 fermion, 3 fermions and 5 fermions and if the action is to be invariant, these must individually cancel.

For example, the one-fermion terms come from  $\delta x^i = \epsilon^\dagger \psi^i$  in terms that started off with no fermions, and from the first part of the fermion variation  $\delta_1 \psi^{\dagger i} = i \epsilon^\dagger \dot{x}^i$  in terms that started off with two fermions. These are

$$\begin{aligned} \delta S \Big|_{1\text{-fermion}} &= \int dt \, g_{ij} \dot{x}^i \delta \dot{x}^j + i g_{ij} \delta_1 \psi^{\dagger i} (\dot{\psi}^j + \Gamma_{kl}^j \dot{x}^k \psi^l) + \frac{1}{2} \delta g_{ij} \dot{x}^i \dot{x}^j \\ &= \int dt \, g_{ij} \dot{x}^i (\epsilon^\dagger \dot{\psi}^j + \epsilon^\dagger \dot{\psi}^j) - g_{ij} \epsilon^\dagger \dot{x}^i (\dot{\psi}^j + \Gamma_{kl}^j \dot{x}^k \psi^l) + \frac{1}{2} \partial_l g_{ij} \epsilon^\dagger \psi^l \dot{x}^i \dot{x}^j \end{aligned}$$

There are two terms with  $\epsilon^\dagger \dot{x} \dot{\psi}$  that immediately cancel. We're left with

$$\delta S \Big|_{1\text{-fermion}} = \int dt \, \epsilon^\dagger \left[ \frac{1}{2} \partial_l g_{ij} - g_{ik} \Gamma_{jl}^k \right] \dot{x}^i \dot{x}^j \psi^{\dagger l} + \epsilon^\dagger g_{ij} \dot{x}^i \dot{\psi}^j$$

The first term vanishes because the metric is covariantly constant,  $\nabla g = 0$ . Or, in more detail, we use the definition of the Levi-Civita connection,  $g_{ik}\Gamma_{jl}^k = \frac{1}{2}(\partial_j g_{il} + \partial_l g_{ij} - \partial_i g_{jl})$ . But this comes multiplied by  $\dot{x}^i \dot{x}^j$  in the variation which means that we get to symmetrise, so  $g_{ik}\Gamma_{jl}^k \dot{x}^i \dot{x}^j = \frac{1}{2}\partial_l g_{ij} \dot{x}^i \dot{x}^j$  which, happily, cancels the other term in the variation of the action. We're left with

$$\delta S \Big|_{1\text{-fermion}} = \int dt \epsilon^\dagger g_{ij} \dot{x}^i \psi^j$$

As explained above, we identify this as the conserved charge arising from the symmetry,  $\delta S = \epsilon^\dagger Q^\dagger$ , giving  $Q^\dagger = g_{ij} \dot{x}^i \psi^j$  as advertised in (3.8).

It's simplest to next look at terms with 5 fermions. These come from the  $\delta R_{ijkl}$  term and the  $R_{ijkl}\psi^i \delta_2 \psi^{\dagger j} \psi^k \psi^{\dagger l}$  term where we include only the part of the fermion variation that itself has two fermions,  $\delta_2 \psi^{\dagger j} = \epsilon^\dagger \Gamma_{mn}^j \psi^{\dagger m} \psi^n$ . Combined, these terms give

$$\delta S \Big|_{5\text{-fermion}} = \int dt \epsilon^\dagger \left[ -\frac{1}{4} \partial_m R_{ijkl} \psi^m \psi^i \psi^j \psi^{\dagger k} \psi^{\dagger l} - \frac{1}{2} R_{ijkl} \Gamma_{mn}^j \psi^i \psi^j \psi^{\dagger m} \psi^n \psi^{\dagger l} \right]$$

After using the fermions to impose anti-symmetry, this term vanishes by virtue of the Bianchi identity  $\nabla_{[m} R_{ij]kl} = 0$ .

This leaves us with the 3-fermions terms in the variation of the action. They are, of course, everything that we didn't yet consider.

$$\begin{aligned} \delta S \Big|_{3\text{-fermion}} &= \int dt \delta g_{ij} \left( i\psi^{\dagger i} \dot{\psi}^j + i\Gamma_{kl}^j \dot{x}^k \psi^{\dagger i} \psi^l \right) + i g_{ij} \delta_2 \psi^{\dagger i} (\dot{\psi}^j + \Gamma_{kl}^j \dot{x}^k \psi^l) \\ &\quad + i g_{ij} \psi^{\dagger i} (\delta \Gamma_{kl}^j \dot{x}^k \psi^l + \Gamma_{kl}^j \delta \dot{x}^k \psi^l) - \frac{1}{2} R_{ijkl} \psi^i \psi^j \delta_1 \psi^{\dagger k} \psi^{\dagger l} \\ &= \int dt \epsilon^\dagger \partial_m g_{ij} \psi^m \left( i\psi^{\dagger i} \dot{\psi}^j + i\Gamma_{kl}^j \dot{x}^k \psi^{\dagger i} \psi^l \right) + i g_{ij} \epsilon^\dagger \Gamma_{mn}^i \psi^{\dagger m} \psi^n (\dot{\psi}^j + \Gamma_{kl}^j \dot{x}^k \psi^l) \\ &\quad - i g_{ij} \epsilon^\dagger \left[ \psi^{\dagger i} \partial_m \Gamma_{kl}^j \psi^m \dot{x}^k \psi^l + \psi^{\dagger i} \Gamma_{kl}^j \dot{\psi}^k \psi^l \right] - \frac{i}{2} \epsilon^\dagger R_{ijkl} \psi^i \psi^j \dot{x}^k \psi^{\dagger l} \end{aligned}$$

There are two different kinds of terms in this expression. The first take the form  $\psi^{\dagger} \dot{\psi} \psi$ . Gathering them together, we find that they come multiplying  $\nabla g = 0$ . The second take the form  $\dot{x} \psi^{\dagger} \psi \psi$ . The first of these involve combinations of the connection that gather together to give  $\partial \Gamma + \Gamma^2$ . But this is the definition of the Riemann tensor and is cancelled by the final term above. The upshot is that, for a global variation with  $\dot{\epsilon}^\dagger = 0$ , we have  $\delta S = 0$ : the action is supersymmetric.

### 3.1.1 Quantisation: Filling in Forms

Quantising the sigma model needs a little care due to operator ordering issues. The canonical momenta are

$$p_i = \frac{\partial L}{\partial \dot{x}^i} = g_{ij} (\dot{x}^j + i\Gamma_{kl}^j \psi^{\dagger k} \psi^l) \quad \text{and} \quad \frac{\partial L}{\partial \psi^i} = -ig_{ij} \psi^{\dagger j}$$

We have, as always

$$[x^i, p_j] = \delta_j^i \quad \text{and} \quad \{\psi^i, \psi^{\dagger j}\} = g^{ij}$$

The tricky commutator is, it turns out, the one between bosons and fermions. This is best described in the terms of the *mechanical momentum* as opposed to the canonical momentum,

$$\pi_i = g_{ij} \dot{x}^j = p_i - ig_{il} \Gamma_{jk}^l \psi^{\dagger j} \psi^k$$

The associated commutation relations turn out to be

$$[\pi_i, \psi^j] = i\Gamma_{ik}^j \psi^k, \quad [\pi_i, \psi^{\dagger j}] = i\Gamma_{ik}^j \psi^{\dagger k} \quad \text{and} \quad [\pi_i, \pi_j] = -R_{ijkl} \psi^{\dagger k} \psi^l$$

Let's now look more closely at the Hilbert space of fermions. We quantise the fermions in the usual way: we introduce a state  $|0\rangle$  that obeys

$$\psi^i |0\rangle = 0$$

for all  $i = 1, \dots, n$ . We then build up the Hilbert space by acting with successive  $\psi^{\dagger i}$ . At the first level we have  $n$  states,  $\psi^{\dagger i} |0\rangle$ . At the next level we have  $\frac{1}{2}n(n-1)$  states,  $\psi^{\dagger i} \psi^{\dagger j} |0\rangle = -\psi^{\dagger j} \psi^{\dagger i} |0\rangle$ , and so on. The natural anti-symmetry of Grassmann objects means that there are  $\binom{n}{p}$  states of the form  $(\psi^{\dagger})^p |0\rangle$ .

As we already advertised in Section 1.4.1, this is a very familiar structure in geometry: it arises for totally anti-symmetric  $(0, p)$  tensor fields, also known as  $p$ -forms. This prompts the identification

$$\begin{aligned} |0\rangle &\longleftrightarrow 1 \\ \psi^{\dagger i} |0\rangle &\longleftrightarrow dx^i \\ \psi^{\dagger i} \psi^{\dagger j} |0\rangle &\longleftrightarrow dx^i \wedge dx^j \\ &\vdots \\ \psi^{\dagger 1} \dots \psi^{\dagger n} |0\rangle &\longleftrightarrow dx^1 \wedge \dots \wedge dx^n \end{aligned}$$

States in the Hilbert space of supersymmetric quantum mechanics are no longer just functions over the manifold  $M$ , but now all forms over the manifold  $M$ . States of the kind  $f(x)(\psi^{\dagger})^p |0\rangle$  correspond to  $p$ -forms. We denote the space of  $p$ -forms over  $M$  as  $\Lambda^p(M)$ .

This relation between Grassmann variables and forms, identifying  $\psi^{\dagger i} \longleftrightarrow dx^i \wedge$  provides the key link between supersymmetry and more interesting aspects of geometry. From this, many lovely geometrical facts follow. For example, we can ask: what is the geometrical interpretation of  $\psi^i$ ? From the commutation relation  $\{\psi^i, \psi^{\dagger j}\} = g^{ij}$ , it clearly acts as a map  $\psi^i : \Lambda^p(M) \mapsto \Lambda^{p-1}(M)$ . We can be more explicit and check

$$\begin{aligned} \psi^i \psi^{\dagger j} \psi^{\dagger k} \dots \psi^{\dagger l} |0\rangle &= \{\psi^i, \psi^{\dagger j} \psi^{\dagger k} \dots \psi^{\dagger l}\} |0\rangle \\ &= [(g^{ij} \psi^{\dagger k} \dots \psi^{\dagger l}) - (\psi^{\dagger j} g^{ik} \dots \psi^{\dagger l}) + \dots] |0\rangle \end{aligned}$$

But, in the language of forms, this is the action of the *interior product*,

$$\psi^i \longleftrightarrow g^{ij} \iota_{\partial/\partial x^j}$$

Meanwhile, the inner product between states in the Hilbert space is,

$$\langle \omega | \eta \rangle = \int_M \bar{\omega} \wedge \star \eta \quad (3.9)$$

Where  $\bar{\omega}$  is the complex conjugation of  $\omega$  and  $\star$  is the Hodge dual. Note that this is non-vanishing only if  $\omega$  and  $\eta$  are forms of the same degree  $p$ . Furthermore, evaluated on functions  $\omega \in \Lambda^0(M)$ , it reproduces the norm (3.5).

The Lagrangian (3.6) has a  $U(1)$  symmetry acting on fermions as

$$\psi^i \rightarrow e^{i\alpha} \psi^i \quad \text{and} \quad \psi^{\dagger i} \rightarrow e^{-i\alpha} \psi^{\dagger i}$$

The corresponding Noether charge is

$$F = g_{ij} \psi^{\dagger i} \psi^j$$

which counts the number of fermionic excitations or, in our new geometrical language, the degree of the form. If we have a state  $|\phi\rangle \in \Lambda^p(M)$ , then

$$F|\phi\rangle = p|\phi\rangle$$

The fact that  $F$  is conserved means that Hamiltonian evolution doesn't mix up forms of different degrees: energy eigenstates lie in a particular  $\Lambda^p(M)$ . The fermion number  $F$  also provides the grading that splits our Hilbert space into bosonic and fermionic pieces:  $\mathcal{H} = \mathcal{H}_B \oplus \mathcal{H}_F$ . These comprise of even and odd forms respectively.

$$\mathcal{H}_B = \mathbb{C} \otimes \left[ \bigoplus_{p \text{ even}} \Lambda^p(M) \right] \quad \text{and} \quad \mathcal{H}_F = \mathbb{C} \otimes \left[ \bigoplus_{p \text{ odd}} \Lambda^p(M) \right]$$

where the overall factor of  $\mathbb{C}$  is there simply because wavefunctions are complex valued in quantum mechanics rather than real.

Finally, we come to the supercharges  $Q$  and  $Q^\dagger$  themselves. The presence of the momentum operator means that these act as derivatives, while the fermions ensure that they also map  $Q : \Lambda^p(M) \mapsto \Lambda^{p+1}(M)$ . But there is a very natural object in differential geometry with these properties: it is the exterior derivative

$$Q = i\psi^\dagger p_i \longleftrightarrow dx^i \wedge \frac{\partial}{\partial x^i} = d$$

Similarly,  $Q^\dagger : \Lambda^p(M) \mapsto \Lambda^{p-1}(M)$  act as the adjoint operator

$$Q^\dagger = i\psi^i p_i \longleftrightarrow g^{ij} \iota_{\partial/\partial x^i} \frac{\partial}{\partial x^j} = d^\dagger$$

Acting on  $p$ -forms, the adjoint operator can also be written as

$$d^\dagger = (-1)^{n(p+1)+1} \star d \star$$

This adjoint operator annihilates functions  $d^\dagger f = 0$  for  $f \in \Lambda^0(M)$ . This is to be expected since it follows from  $\psi^i|0\rangle = 0$ . Similarly, the exterior derivative itself annihilates top forms,  $d\omega = 0$  for all  $\omega \in \Lambda^n(M)$ . Before we go on, note that the correspondence  $Q \equiv d$  and  $Q^\dagger \equiv d^\dagger$  is the reason that we chose to define  $Q$  to be the supercharge involving  $\psi^\dagger$  rather than, as in Section 1, in terms of  $\psi$ .

The identification of the supercharges also gives a geometric meaning to the Hamiltonian. It is

$$H = \frac{1}{2}\{Q, Q^\dagger\} \quad \Rightarrow \quad H = \frac{1}{2}\Delta$$


with

$$\Delta = dd^\dagger + d^\dagger d$$

This is the *Laplacian* operator in differential geometry. It is clear from its definition in terms of  $d$  and  $d^\dagger$  that it is a prime candidate for a supersymmetric Hamiltonian; in some sense everything that we've done above is just to realise this possibility in terms of Grassmann variables  $\psi$  and  $\psi^\dagger$ .

The Laplacian is positive definite, as befits a supersymmetric Hamiltonian. This follows from the fact that the  $\dagger$  in  $Q^\dagger$  (or, equivalently  $d^\dagger$ ) means the adjoint operation with respect to the inner product (3.9) so that, for any  $\omega \in \Lambda^p(M)$ ,

$$\langle \omega | \Delta \omega \rangle = \langle \omega | dd^\dagger \omega \rangle + \langle \omega | d^\dagger d \omega \rangle = \|d^\dagger \omega\|^2 + \|d\omega\|^2 \geq 0 \quad (3.10)$$



**Figure 8.** A typical energy spectrum of the quantum mechanics, with pairs of states related by supersymmetry in yellow and the lone ground states, associated to de Rham cohomology, in red.

A short calculation (see, for example, Section 3.1.4 of the lectures on [General Relativity](#)) shows that, when acting on function  $f \in \Lambda^0(M)$ , the Laplacian is given by

$$\Delta f = \frac{1}{\sqrt{g}} \partial_i (\sqrt{g} g^{ij} \partial_j f)$$

in agreement with the Hamiltonian (3.3) for the non-supersymmetric sigma model. Note, however, that in the absence of supersymmetry there was always the option to add the  $\alpha R$  term in (3.3) to the Hamiltonian. Supersymmetry removes this ambiguity.

### 3.1.2 Ground States and de Rham Cohomology

We'll now consider the kind of spectrum that we expect to find. As we saw in Section 1, all states with energy  $E \neq 0$  must come in pairs. In particular, if an energy eigenstate with  $E \neq 0$  obeys

$$Q|\alpha\rangle = 0$$

then  $|\alpha\rangle$  is *Q-exact*, meaning that it can be written as  $|\alpha\rangle = Q|\phi\rangle$  for some  $|\phi\rangle$ . To see this, we just need to use  $QQ^\dagger + Q^\dagger Q = 2E$  to see that

$$|\alpha\rangle = \frac{1}{2E} Q(Q^\dagger|\alpha\rangle)$$

This tells us that  $Q$  and  $Q^\dagger$  map us back and forth between the two states related by supersymmetry. In the form language, we see that supersymmetry relates pairs of  $p$  and  $p + 1$  forms. These are shown as the yellow dots in Figure 8.

However, as we've seen in previous examples, the ground states with  $E = 0$  are special since there is no need for these to be paired. In the present context, the ground states arise from forms that obey

$$\Delta\gamma = 0 \quad \Leftrightarrow \quad d\gamma = d^\dagger\gamma = 0$$

Forms of this kind are called *harmonic*. These are depicted as red dots in Figure 8. The space of harmonic  $p$ -forms is denoted  $\text{Harm}^p(M)$ . We learn that the Hilbert space of ground states is

$$\mathcal{H}_{\text{ground}} = \bigoplus_p \text{Harm}^p(M)$$

This discussion also tells us that there are three kinds of states in the Hilbert space: those for which  $|\phi\rangle = Q|\alpha\rangle$  or  $|\phi\rangle = Q^\dagger|\beta\rangle$ , which sit in supersymmetric pairs. And those for which  $Q|\phi\rangle = Q^\dagger|\phi\rangle = 0$  which are the supersymmetric ground states. This means that any state  $|\phi\rangle \in \mathcal{H}$  has a unique decomposition as

$$|\phi\rangle = Q|\alpha\rangle + Q^\dagger|\beta\rangle + |\omega\rangle \tag{3.11}$$

where  $\Delta|\omega\rangle = 0$ . In the geometric language, this is equivalent to saying that any form can be written uniquely as

$$\omega = d\alpha + d^\dagger\beta + \gamma \tag{3.12}$$

where  $\omega$  is harmonic. This is known as the *Hodge decomposition theorem*.

There is an important comment to make here. The Hodge decomposition theorem is not a trivial statement in mathematics. It took Hodge much of the 1930s to prove and, even then, needed corrections from Weyl and Kodaira. Yet the statement about the decomposition of states in the Hilbert space (3.11) follows trivially from the structure of supersymmetric quantum mechanics! What's going on?

Shortly we will “prove” other theorems in geometry where we will make use of the physicist's secret weapon, the path integral. Here, however, the power of physics comes only from our blatant disregard for anything approaching rigour. In geometry, the space of differential forms is *not* a Hilbert space because the inner product (3.9) is not complete. In quantum mechanics, we deal with this by restricting attention to  $L^2$  forms but then one has to worry whether the exterior derivative acts solely within this space. All of these are subtleties that we sweep under the rug in physics, but present the real challenge behind the proof of the Hodge decomposition theorem.

## Cohomology

There is another way to view the ground states in terms of *cohomology*. As we've seen, the exterior derivative  $d$  (or equivalently the supercharge  $Q$ ) maps us from  $d : \Lambda^p(M) \rightarrow \Lambda^{p+1}(M)$ . We can depict this in terms of what mathematicians call a *chain complex*

$$\Lambda^0(M) \xrightarrow{d} \Lambda^1(M) \xrightarrow{d} \Lambda^2(M) \xrightarrow{d} \Lambda^3(M) \xrightarrow{d} \dots$$

Because  $d^2 = 0$ , the image of one map necessarily lies in the kernel of the next. The idea of cohomology is that it's interesting to look more closely at the difference between the kernel and image.

First some definitions. A form  $\phi$  is said to be *closed* if  $d\phi = 0$ . We denote the space of all closed  $p$ -forms as  $Z^p(M)$ . Another way to say this is that  $Z^p(M)$  is the kernel of the map  $d : \Lambda^p(M) \rightarrow \Lambda^{p+1}(M)$ .

A form  $\phi$  is said to be *exact* if it can be written as  $\phi = d\alpha$  for some  $\alpha$ . We denote the space of all exact  $p$ -forms as  $B^p(M)$ . Another way to say this is that  $B^p(M)$  is image of the map  $d : \Lambda^{p-1}(M) \rightarrow \Lambda^p(M)$ .

As we mentioned above, we necessarily have  $B^p(M) \subset Z^p(M)$ . The *de Rham cohomology group* is defined to be

$$H^p(M) = Z^p(M)/B^p(M)$$

The quotient here is an equivalence class. Two closed forms  $\phi$  and  $\phi' \in Z^p(M)$  are said to be equivalent if  $\phi = \phi' + d\alpha$  for some  $\alpha$ . We say that  $\phi$  and  $\phi'$  sit in the same equivalence class  $[\phi]$ . The cohomology group  $H^p(M)$  is the set of equivalence classes. In other words, it consists of closed forms mod exact forms.

Finally, we define the *Betti numbers*,

$$b_p = \dim H^p(M)$$

There are a number of interesting things about these Betti numbers. First, this counting of cohomology classes is just another way of counting the ground states in quantum mechanics, and the Betti numbers can equally well be viewed as counting harmonic forms. This follows from...

**Claim:** There is an isomorphism  $H^p(M) \cong \text{Harm}^p(M)$  and so

$$b_p = \dim \text{Harm}^p(M)$$

**Proof:** The proof follows straightforwardly from the Hodge decomposition (3.12). We'll first show that each harmonic form is associated to an element of  $H^p(M)$ . Clearly any harmonic form  $\gamma$  is closed, with  $d\gamma = 0$ . But the unique nature of the Hodge decomposition (3.12) means that  $\gamma$  cannot be written as  $\gamma = d(\text{something})$  and so forms the basis of an equivalence class  $[\gamma] \in H^p(M)$ .

Next we must go the other way and show that each equivalence class of  $[\omega] \in H^p(M)$  is associated to a harmonic form. We decompose  $\omega = d\alpha + d^\dagger\beta + \gamma$ . By the definition of  $[\omega] \in H^p(M)$ , we must have  $d\omega = 0$  and so, using the inner product (3.9), we have


$$0 = \langle d\omega | \beta \rangle = \langle \omega | d^\dagger\beta \rangle = \langle d\alpha + d^\dagger\beta + \gamma | d^\dagger\beta \rangle = \langle d^\dagger\beta | d^\dagger\beta \rangle$$

where, in the final step, we integrated by parts and used the facts that  $dd\alpha = 0$  and  $d\gamma = 0$ . The upshot is that  $d^\dagger\beta = 0$  and any element of the equivalence class  $[\omega] \in H^p(M)$  takes the form  $\omega = d\alpha + \gamma$ . Any other member of the same equivalence class  $\omega' \in [\omega]$  can be written as  $\omega' = d\eta + \gamma$  and is associated to the same harmonic form  $\gamma$ .  $\square$

There's an analogy here with gauge symmetry that is worth highlighting. In Maxwell theory, the gauge potentials  $A$  and  $A + d\alpha$  are physically equivalent as they are related by a gauge transformation. If we want to pick a representative of this equivalence class then we need gauge fixing condition that picks out one particular choice of  $A$ . For cohomology, the equivalence class  $[\omega]$  relates  $\omega \sim \omega + d\alpha$ . A representative of this class can be picked by the "gauge fixing condition"  $d^\dagger\omega = 0$ . This then picks out the harmonic forms as special.

Any manifold  $M$  with dimension  $\dim(M) = n$  always has  $b_0 = 1$  and  $b_n = 1$ . The zero forms are just functions over the manifold, and any constant function over  $M$  is clearly harmonic, but cannot be written as  $d(\text{something})$  as there are no  $p = -1$  forms. Similarly, the volume form  $\text{Vol} = \star 1$  provides the harmonic top form.

Other Betti numbers come in pairs with  $b_p = b_{n-p}$ , a relationship that follows from Poincaré duality. It turns out that all these higher Betti numbers are non-vanishing only if the manifold  $M$  has some interesting topology. To explain this, we need to remove the co in cohomology.



**Figure 9.** The red lines depict a topologically trivial submanifold  $C$  on the left, and a topologically non-trivial sub-manifold  $C$  on the right.

## Homology

Here we give a brief overview of how the de Rham cohomology, and associated harmonic forms, contain information about the topology of the manifold  $M$ .


Consider a submanifold  $C \subset M$ . We'll take this to be a closed submanifold, meaning that it has no boundary

$$\partial C = 0$$

An interesting question is whether  $C$  itself can be thought of as the boundary of another manifold, meaning  $C = \partial D$ . This is a question of topology.

We can see this in two simple examples shown in Figure 9. There we depict two manifolds of dimension two: the sphere  $M = \mathbf{S}^2$  and the torus  $M = \mathbf{T}^2$ . On each we've drawn a one-dimensional submanifold  $C$  as a red line. For  $C \subset \mathbf{S}^2$ , this submanifold is the boundary of a disc  $C = \partial D$ . For  $C \subset \mathbf{T}^2$  there is no such bounding manifold  $D$ . This reflects the fact that there is interesting topology in the torus, but not in the sphere.

Indeed, there are actually two different topologically non-trivial submanifolds of the torus: in addition to the circle  $C$  shown in Figure 9, there is also the circle  $C'$  that winds in the way shown on the right.



The algebraic structure of these topologically non-trivial submanifolds is identical to those of forms. In particular, a boundary of a boundary is always vanishing, which we write as  $\partial^2 = 0$ . This, obviously, is the

strikingly reminiscent of the exterior derivative relation  $d^2 = 0$ . We use this to define homology groups using  $\partial$  analogous to the cohomology groups that we defined previously using  $d$ . The *homology group*  $H_p(M)$  is the equivalence class of closed  $p$ -dimensional submanifolds that are not themselves the boundary of a  $(p+1)$ -dimensional manifold. In particular, two submanifolds  $C_1$  and  $C_2$  lie in the same cohomology class if one can be smoothly deformed into the other. In terms of equations, this means that difference is a boundary,

$$C_1 \sim C_2 \text{ if and only if } C_1 - C_2 = \partial D$$

The relationship between homology and cohomology is more than just an analogy. The spaces  $H_p(M)$  and  $H^p(M)$  are dual to each other, and hence isomorphic. This statement, known as *de Rham's theorem*, is not straightforward to prove but it's easy to get some intuition for how it works. Given a closed submanifold  $C \subset M$  and a form  $\omega$  on  $M$  we can define a map to the real numbers, given by

$$(C, \omega) = \int_C \omega$$

Strictly speaking, the integral only makes sense if  $\dim C = p$  and  $\omega$  is a  $p$ -form. If the form  $\omega$  has a degree different than  $\dim C$  then the pairing is simply said to be zero. In what follows, we will sometimes refer to such a closed submanifold  $C$  as a *cycle*.

This pairing has some lovely properties that follow from Stokes' theorem. First, the answer depends only on the equivalence class  $[\omega] \in H^p(M)$ . To see this, note that

$$(C, \omega + d\alpha) = \int_C (\omega + d\alpha) = \int_C \omega + \int_C d\alpha$$

but the total derivative  $\int_C d\alpha = 0$  because  $\partial C = 0$ .

Conversely, if we consider two submanifolds  $C_1$  and  $C_2$  that can be smoothly deformed into each other, so  $C_1 - C_2 = \partial D$ , then integrating any closed form  $\omega$  gives


$$\int_{C_1} \omega - \int_{C_2} \omega = \int_{\partial D} \omega = \int_D d\omega = 0$$

We see that the answer only depends on the equivalence class  $[C] \in H_p(M)$ .

The upshot of these arguments is that the ground states of the supersymmetric sigma model (3.6) are determined by the topology of the target space  $M$ . Heuristically, the quantum particle can minimise its energy by spreading its wavefunction over topologically non-trivial submanifolds of  $M$ .

Here are some simple examples. The sphere  $\mathbf{S}^n$  is boring: its only Betti numbers are  $b_0 = 1$  corresponding to constant functions and  $b_n = 1$  corresponding to the top form. The torus  $\mathbf{T}^n$  is more interesting: it has Betti numbers  $b_p = \binom{n}{p}$ .

In  $n = 2$  dimensions, closed manifolds are known as Riemann surfaces and are labelled by their genus  $g$  which counts the number of holes. Here are some examples of manifolds with genus  $g = 0$ ,  $g = 1$  and  $g = 2$  respectively



The Betti numbers are  $b_0 = b_2 = 1$  and  $b_1 = 2g$ . Each extra hole introduces two new topologically non-trivial 1-manifolds that encircle the hole in different ways.

Finally, I should mention in any logical presentation, homology precedes cohomology. Our physics approach has lead us to introduce these in an inverted order.

### 3.1.3 The Witten Index and the Chern-Gauss-Bonnet Theorem

In Section 1, we learned that there is something special about the Witten index in supersymmetric quantum mechanics. Recall that this is defined by  $\text{Tr} (-1)^F e^{-\beta H}$  and counts then number of supersymmetric ground states, up to a sign.

For our supersymmetric sigma model, the Witten index is just the alternating sum of Betti numbers

$$\text{Tr} (-1)^F e^{-\beta H} = \chi(M) := \sum_p (-1)^p b_p \quad (3.13)$$

This is perhaps the most famous topological invariant in mathematics: it is known as the *Euler character* of the manifold.

Again, some examples. The sphere  $\mathbf{S}^n$  has Euler character

$$\chi(\mathbf{S}^n) = 1 + (-1)^n$$

so is either  $\chi(\mathbf{S}^n) = 2$  for  $n$  even or  $\chi(\mathbf{S}^n) = 0$  for  $n$  odd. The torus  $\mathbf{T}^n$  always has  $\chi(\mathbf{T}^n) = 0$ . The 2d Riemann surface of genus  $g$  has  $\chi(M) = 2 - 2g$ .

We can see from our discussion of quantum mechanics why this is a topological invariant. We know that the Witten index is robust against any small change of the parameters in the quantum mechanics. In the present case, that means that if we vary the metric  $g_{ij}$ , at least within reason so that we avoid singularities, then the Witten index should remain unchanged. But that means that object  $\chi(M)$  defined in (3.13) must be independent of the the choice of metric: it can be depend only on cruder aspects of  $M$ , specifically its topology.

Finally, note that the sigma models provide many other examples in which the Witten index vanishes but there are, nonetheless, ground states with  $E = 0$ . For example, the sigma model on  $\mathbf{S}^3$  (or, indeed, any odd dimensional sphere) has  $\chi(\mathbf{S}^3) = 0$  but there are two ground states, one the constant function corresponding  $b_0 = 1$  and the other the volume form corresponding to  $b_3 = 1$ . These ground states are also protected by topology, this time by the cohomology rather than the cruder Euler character.

### The Path Integral Again

As we saw in Section 2, there is a straightforward description of the Witten index in terms of the path integral. We simply need to calculate

$$\mathcal{I} = \text{Tr} (-1)^F e^{-\beta H} = \int \mathcal{D}x \mathcal{D}\psi^\dagger \mathcal{D}\psi e^{-S_E[x, \psi, \psi^\dagger]}$$

where Euclidean time  $\tau$  has period  $\beta$  and both  $x$  and  $\psi$  are assigned periodic boundary conditions. The Euclidean action is

$$S_E = \oint d\tau \frac{1}{2} g_{ij}(x) \dot{x}^i \dot{x}^j + g_{ij} \psi^{\dagger i} \nabla_\tau \psi^i + \frac{1}{4} R_{ijkl} \psi^i \psi^j \psi^{\dagger k} \psi^{\dagger l}$$

with  $\nabla_\tau \psi^i = \dot{\psi}^i + \Gamma_{jk}^i \dot{x}^j \psi^k$ . We know that the Witten index is independent of  $\beta$ . We will use this to compute the path integral in the limit  $\beta \rightarrow 0$ . The key idea is that, in this limit, any non-trivial excitations around the Euclidean circle costs an increasing amount of action and so we can restrict ourselves to constant configurations, where the path integral reduces to a normal integral.

Putting these words into formulae, we first rescale the time coordinate to work with  $\tau' = \tau/\beta$  so the new time coordinate has period  $\tau' \in [0, 1)$ . We also rescale  $\psi \rightarrow \beta^{-1/4} \psi$ , leaving us with the Euclidean action

$$S_E = \int_0^1 d\tau' \frac{1}{2\beta} g_{ij}(x) \dot{x}^i \dot{x}^j + \frac{1}{\sqrt{\beta}} g_{ij} \psi^{\dagger i} \nabla_\tau \psi^i + \frac{1}{4} R_{ijkl} \psi^i \psi^j \psi^{\dagger k} \psi^{\dagger l}$$

where we now see explicitly that in the limit  $\beta \rightarrow 0$ , the modes with  $\dot{x}$  and  $\dot{\psi}$  non-zero are heavily suppressed. The path integral then reduces to the ordinary integral

$$\text{Tr} (-1)^F e^{-\beta H} = \frac{1}{(2\pi)^{n/2}} \int d^n x \frac{1}{\sqrt{g}} \int d^n \psi d^n \psi^\dagger \exp \left( -\frac{1}{4} R_{ijkl} \psi^i \psi^j \psi^{\dagger k} \psi^{\dagger l} \right)$$

As in previous examples, we have to saturate the Grassmann integration. But this time, there's clear way to do it. We simply expand out the exponential until we find the right number of fermions.

Since the fermions always come in groups of four, if  $n$  is odd the integral necessarily vanishes. We learn that

$$\chi(M) = 0 \quad \text{if } \dim M = \text{odd}$$

This simple result also follows from the relation  $b_p = b_{n-p}$ . However, if  $n$  is even then the term with  $n/2$  powers of the Riemann tensor will saturate the integral.

We start with  $n = 2$ . In this case, we pull down just a single copy of the Riemann tensor. After doing the Grassmann integrations, we find

$$\text{Tr} (-1)^F e^{-\beta H} = \frac{1}{4\pi} \int d^2 x \sqrt{g} R$$

This is the well known Gauss-Bonnet expression for the Euler character of a Riemann surface.

In general, the Grassmann integrations leave us with  $n/2$  copies of the Riemann tensor, contracted with epsilon symbols

$$\text{Tr} (-1)^F e^{-\beta H} = \frac{1}{(4\pi)^{n/2} (n/2)!} \int d^n x \frac{1}{\sqrt{g}} \epsilon^{i_1 \dots i_n} \epsilon^{j_1 \dots j_n} R_{i_1 i_2 j_1 j_2} \dots R_{i_{n-1} i_n j_{n-1} j_n}$$

This is the generalisation of the Gauss-Bonnet theorem, first proven by Chern in 1944. The contraction of the epsilon symbols results in an expression known as the *Euler density*. The slightly unusual looking  $1/\sqrt{g}$  should be thought of as  $\sqrt{g} \times \frac{1}{\sqrt{g}} \times \frac{1}{\sqrt{g}}$  with the  $\frac{1}{\sqrt{g}}$  factors combining with the epsilon symbols to give tensor densities.

As an example, for  $n = 4$  dimensional manifolds the expansion of the Euler density gives

$$\chi(M) = \frac{1}{8\pi^2} \int_M d^4 x \sqrt{g} (R_{ijkl} R^{ijkl} - 4R_{ij} R^{ij} + R^2)$$

The magic of the Chern-Gauss-Bonnet theorem is that a global topological object,  $\chi(M)$ , is described in terms of an integral of local data, the Euler density. The magic of supersymmetric quantum mechanics is that it gives a straightforward derivation of this result, with the only real complication the combinatoric factors that arise from Grassmann integration. This is first example where a deep mathematical result can be derived in a different way using the path integral. It won't be the last.

### 3.2 Morse Theory

Our goal in this section is to understand some basic ideas of Morse theory, viewed through the lens of supersymmetric quantum mechanics.

We stick with our  $N = 2$  supersymmetric sigma model (3.6), describing a particle moving on a manifold  $M$ . The novelty is that we now also include a potential  $h(x)$  over the manifold. The resulting supersymmetric theory is a combination of the sigma model and the kind of theories we considered in Section 1.4.1,

$$L = \frac{1}{2} g_{ij} \dot{x}^i \dot{x}^j + i g_{ij} \psi^{\dagger i} \nabla_t \psi^j - \frac{1}{4} R_{ijkl} \psi^i \psi^j \psi^{\dagger k} \psi^{\dagger l} - \frac{1}{2} g^{ij} \frac{\partial h}{\partial x^i} \frac{\partial h}{\partial x^j} - (\nabla_i \partial_j h) \psi^{\dagger i} \psi^j \quad (3.14)$$

Note that the final, fermionic term has the opposite sign from that of Section 1.3; this is purely a choice of convention and, as we will see shortly, will bring us in line with definitions used in mathematics. This action is invariant under the supersymmetry transformations

$$\begin{aligned} \delta x^i &= \epsilon^\dagger \psi^i - \epsilon \psi^{\dagger i} \\ \delta \psi^i &= \epsilon \left( -i \dot{x}^i + \Gamma_{jk}^i \psi^{\dagger j} \psi^k - g^{ij} \frac{\partial h}{\partial x^j} \right) \\ \delta \psi^{\dagger i} &= \epsilon^\dagger \left( +i \dot{x}^i + \Gamma_{jk}^i \psi^{\dagger j} \psi^k - g^{ij} \frac{\partial h}{\partial x^j} \right) \end{aligned} \quad (3.15)$$

These are a combination of the transformations (1.19) for our original quantum mechanics with a potential and (3.7) for the supersymmetric sigma model.

In the absence of the potential, we know that the ground states of the supersymmetric quantum mechanics spread over cycles of  $M$ . However, when we add a the potential  $h$ , the wavefunctions get squeezed and, as the potential gets larger, the wavefunctions are increasingly localised at the minima of the potential. We know that the Witten index can't change. But, more strongly, the total number of  $E = 0$  ground states doesn't change either and, even in the presence of the potential, is given by the Betti numbers of the manifold.

To see this statement, first note that the supercharges associated to (3.15) are given by

$$Q = \left( g_{ij} \dot{x}^i + i \frac{\partial h}{\partial x^j} \right) \psi^{\dagger j} \quad \text{and} \quad Q^\dagger = \left( g_{ij} \dot{x}^i - i \frac{\partial h}{\partial x^j} \right) \psi^j$$

Translated into the geometric language, we have

$$Q \longleftrightarrow d + dx^i \wedge \partial_i h = d + dh \wedge = e^{-h} de^h \quad (3.16)$$

Similarly

$$Q^\dagger \longleftrightarrow (d + dh \wedge)^\dagger = e^h d^\dagger e^{-h}$$

We saw in Section 3.1.2 that the ground states are determined by the cohomology of  $Q$ . But the cohomology when  $h \neq 0$  is isomorphic to the cohomology when  $h = 0$ . We simply take the wavefunctions in the latter case and multiply them by  $e^{-h}$ . Indeed, this is the form of the wavefunctions (1.12) that we found back Section 1.2 when considering a particle on a line.

The fact that the number of supersymmetric ground states is independent of  $h$  means that something interesting must be going on. Because if we crank up  $h$  to be very large, the ground states are localised around the minima of the potential  $V = |\partial_i h|^2$ . This means that there must be some relationship between these minima and the topology of the manifold. This relationship goes under the name of *Morse theory*.

The minima lie at critical points of  $h$  which we will label  $x = X$ . They obey

$$\frac{\partial h}{\partial x^i}(X) = 0 \quad \text{for all } i = 1, \dots, n$$

The function  $h$  is said to be a *Morse function* if it has the property that the critical points  $x = X$  are isolated and non-degenerate. From now on, we'll assume that this is the case.

Consider the situation where we scale the Morse function  $h(x) \rightarrow \zeta h(x)$ , and subsequently send  $\zeta \rightarrow \infty$ . In this limit, the physics is entirely dominated by the critical points of the potential and, at the semi-classical level, the ground state wavefunction is localised at the critical point  $x = X$ . That's not to say that all critical points are necessarily true  $E = 0$  ground states; there may well be tunnelling of the kind that we discussed in Section 2.3 that lifts putative ground states in pairs. But the true ground states must be contained within the set of critical points.

We also need to figure out what's going on with fermions. This is the same calculation that we already met in Section 1.4.1. There, we learned that we should look at the eigenvalues of the Hessian  $\partial_i\partial_j h$ ,

$$(\partial_i\partial_j h) e_k^j = \lambda_k e_k^j$$

where  $e_k^j$  are the eigenvectors and  $\lambda_k$  the eigenvalues, with  $k = 1, \dots, n$ . (The index  $k$  labels the eigenvectors and eigenvalues and shouldn't be summed over. Note also that we flipped the sign of  $h$  in the action (3.14) relative to our discussion in Section 1.4.1, and that shows up as a change of minus sign in this equation relative to (1.27).)

For each negative eigenvalue  $\lambda_k < 0$ , the final term in (3.14) tells us that we can lower the energy by exciting the corresponding collection of fermions  $e_k^j \psi^{\dagger j}$ . Meanwhile, for each positive eigenvalue  $\lambda_k > 0$ , we're better off in the unexcited state.

We define the *Morse index*,  $\mu(X)$ , to be

$$\mu(X) = \text{The number of negative eigenvalues of } \partial_i\partial_j h(X)$$

We learn that the semi-classical ground state sits in the sector with  $\mu(X)$  fermions excited. In other words, the semi-classical ground state at the critical point  $x = X$  is a  $p$ -form with  $p = \mu(X)$ .

Already we learn something striking. We can compute the Witten index by simply summing over the critical points  $X$ , just as we did in (1.24). The novelty is that we know that, for our supersymmetric sigma model, the Witten index tells us the Euler character of the manifold  $M$ . This means that we can compute the Euler character of  $M$  from the critical points of a function over  $M$ ,


$$\chi(M) = \sum_X (-1)^{\mu(X)}$$

In fact, we can say more than this. The total number of critical points may well be more than the total number of  $E = 0$  ground states, since states can be lifted in pairs. But the number of critical points can never be smaller than the number of ground states! Suppose that there are  $m_p$  critical points  $X$  with Morse index  $p = \mu(X)$ . This can be no less than the number of ground states associated to  $p$ -forms, so

$$m_p \geq b_p \tag{3.17}$$

with  $b_p$  the Betti number. This is known as the *weak Morse inequalities*.

Nice as this is, it's possible to do better. We can, in fact, recover the original Betti numbers  $b_p$  from an understanding of the critical points and the relationships between them. In the rest of this section we explain how.



**Figure 10.** Two shapes, both topologically  $\mathbf{S}^2$  with the Morse function given by the height. On the left there are two critical points, on the right there are four. My wife thought it important to point out that these are not drawn to scale.

### A Simple Example: The Two Sphere

To illustrate these ideas, we can look at the case of  $\mathbf{S}^2$ . We know that the Betti numbers are  $b_0 = b_2 = 1$  and  $b_1 = 0$ .

Suppose that we embed  $\mathbf{S}^2$  with its round metric in  $\mathbb{R}^3$ . Then we can consider the height function

$$h = z$$

This is shown in the left-hand side of Figure 10. Clearly there are two critical points of the height function: at the bottom of the sphere where it is a minimum and at the top of the sphere where it is a maximum. The Morse index is  $\mu = 0$  and  $\mu = 2$  respectively, so from the discussion above we know that these ground states are associated to 0-forms and 2-forms. We also know that ground states localised around these minima must be exact  $E = 0$  states.

Now we deform the system. We could change the Morse function  $h$  but, for illustrative purposes, it is simplest if we instead change the metric on the sphere. We'll turn it into the bean shape shown in the right-hand side of Figure 10, keeping the same height function  $h = z$ . This time there are four critical points, one with  $\mu = 0$ , two at the top with  $\mu = 2$ , and the saddle point in the middle with  $\mu = 1$ .

Note that the Euler character hasn't changed,

$$\chi(\mathbf{S}^2) = \sum_X (-1)^{\mu(X)} = 1 + (-1) + 1 + 1 = 2$$

Moreover, the weak Morse inequalities hold, with  $m_0 = b_0 = 1$  and  $1 = m_1 > b_1 = 0$  and  $2 = m_2 > b_2 = 1$ .

For the bean shaped metric, we know that two of four semi-classical ground states must be lifted to have  $E > 0$ . Clearly, it should be the 1-form and some combination of the two 2-forms that gets lifted. Our goal now is to understand how this works, both in the case of the kidney bean and more generally. We will see that much of the technology that we will need has already been covered in the supersymmetric instanton calculation of Section 2.3

### 3.2.1 Instantons Again

Suppose that our Morse function has  $r$  critical points at  $x = X_a$  with  $a = 1, \dots, r$ , such that

$$\frac{\partial h}{\partial x^i}(X_a) = 0$$

The weak Morse inequalities (3.17) tell us that  $r \geq \sum_p b_p$ , the total number of supersymmetric ground states (counted without sign). If  $r = \sum_p b_p$  then the Morse inequalities are saturated,  $m_p = b_p$ , and we're done: as we crank up the strength of the potential, the ground state wavefunctions morph smoothly from being spread over cycles of the manifold, to being localised at the critical points. This is the situation depicted by the orange in the previous example.

However if  $r > \sum_p b_p$ , like in the example of the kidney bean, then there are more critical points than genuine ground states and we have some work to do. Some of the semi-classical ground states associated to critical points must be lifted.

The exact energy eigenstate localised around  $x = X_a$  will be denoted as  $|\Psi_a\rangle$ . Some of these states will persist as zero energy states when all quantum corrections are taken into account. Meanwhile others will be lifted but, as we saw in Section 2.3, will remain as low lying states, with energies of order  $e^{-S_{\text{inst}}}$ . Our goal is to understand this spectrum.

To this end, we will compute the matrix elements

$$\langle \Psi_a | Q | \Psi_b \rangle$$

Any state with  $E = 0$  must be annihilated by  $Q$  so, in general, we expect this matrix to have rank  $r - \sum_p b_p$ , with the zero eigenvectors the true quantum ground states and the remainder those that are lifted.

As in Section 2.3, it's simpler to compute the related set of matrix elements

$$\langle \Psi_a | Q | \Psi_b \rangle \approx \frac{\langle \Psi_a | [Q, h] | \Psi_b \rangle}{h(X_b) - h(X_a)}$$

where, after Wick rotation, the commutator with the supercharge (3.16) gives

$$[Q, h(x)] = \frac{\partial h}{\partial x^i} \psi^{\dagger i}$$

The fact that we have just a single fermion  $\psi^{\dagger}$  in the matrix element means that we'll get non-vanishing contributions if the state  $|\Psi_a\rangle$  has one additional fermion excited than  $|\Psi_b\rangle$ . Or, said differently, if the Morse indices differ by one:

$$\mu(X_a) - \mu(X_b) = 1 \tag{3.18}$$

The difference  $\Delta\mu = \mu(X_a) - \mu(X_b)$  is called the *relative Morse index*.

### The Instanton Equations

It's clear that we are now back in the realm of the quantum tunnelling calculations that we performed in Sections 2.2 and 2.3. To start, we can study the instantons in a sigma model with potential. Focussing just on the bosonic fields for now, the action (3.14) in Euclidean time is

$$S_E = \int d\tau \frac{1}{2} g_{ij} \dot{x}^i \dot{x}^j + \frac{1}{2} g^{ij} \frac{\partial h}{\partial x^i} \frac{\partial h}{\partial x^j}$$

where now  $\dot{x} = dx/d\tau$ . We can write this by completing the square as

$$S_E = \int d\tau \frac{1}{2} g_{ik} \left( \frac{dx^i}{d\tau} \mp g^{ij} \frac{\partial h}{\partial x^j} \right) \left( \frac{dx^k}{d\tau} \mp g^{kl} \frac{\partial h}{\partial x^l} \right) \pm \frac{dx^i}{d\tau} \frac{\partial h}{\partial x^i}$$

For the class of configurations that interpolate from  $x(\tau) = X_b$  at time  $\tau \rightarrow -\infty$  to  $x(\tau) = X_a$  at time  $\tau \rightarrow +\infty$ , the action is minimised for configurations that obey the instanton equations

$$\frac{dx^i}{d\tau} = \pm g^{ij} \frac{\partial h}{\partial x^j}$$

Solutions to the equation with the + sign are instantons; those with the – sign are anti-instantons. The action is given by

$$S_{\text{inst}} = \pm(h(X_a) - h(X_b))$$

Since we want the action to be positive definite, we should pick the instanton solution when  $h(X_a) > h(X_b)$  and the anti-instanton when  $h(X_a) < h(X_b)$ .

From our previous calculation in Section 2.3, we know that the fermion zero modes play a crucial role in supersymmetric instanton calculations. So our next question: how many fermi zero modes does our instanton have? To answer this, we look at the linearised fermion equation of motion. Here “linearised” means that we drop the Riemann tensor term in (3.14), and the connection term in  $\nabla_t$ . In Euclidean time, the linearised equations are

$$D\psi^i := \frac{d\psi^i}{d\tau} + g^{ij}\nabla_j\partial_k h \psi^k = 0 \quad (3.19)$$

and

$$D^\dagger\psi^{\dagger i} := -\frac{d\psi^{\dagger i}}{d\tau} + g^{ij}\nabla_j\partial_k h \psi^{\dagger k} = 0$$

We want to know how many solutions each of these equations have in the background of an instanton. In fact, we really just want to know the difference between the number of solutions to these equations. This is because if both  $D$  and  $D^\dagger$  have zero modes then they will most likely be lifted by the non-linear terms in the action. And, indeed, generically, this will happen. However if there are unpaired zero modes of, say  $D^\dagger$ , then these must be saturated in some other way in the path integral. This prompts our interest in the index

$$\mathcal{I}(D) = \dim \text{Ker } D - \dim \text{Ker } D^\dagger$$

where  $\text{Ker } D$  is the kernel of  $D$ , the space of solutions to  $D\psi^i = 0$ . Clearly, the index counts the number of unpaired zero modes of the instanton.

Furthermore, because our matrix element  $\langle \Psi_a | [Q, h] | \Psi_b \rangle$  contains just a single fermion  $\psi^\dagger$ , we’re only going to get non-zero contributions from instantons which have one more zero mode for  $\psi^\dagger$  than for  $\psi$ , namely

$$\mathcal{I}(D) = -1 \quad \text{or, equivalently} \quad \mathcal{I}(D^\dagger) = +1$$

We’re always guaranteed to get some fermi zero modes from acting with broken supersymmetry. If we start from an instanton configuration, and originally set all fermions

to zero, then acting with the supersymmetry transformations (3.15) (and remembering to Wick rotate to Euclidean time  $\tau = it$ ), we have

$$\begin{aligned}\delta\psi^i &= \epsilon \left( \frac{dx^i}{d\tau} - g^{ij} \frac{\partial h}{\partial x^j} \right) \\ \delta\psi^{\dagger i} &= -\epsilon^\dagger \left( \frac{dx^i}{d\tau} + g^{ij} \frac{\partial h}{\partial x^j} \right)\end{aligned}$$

Because we want a  $\psi^\dagger$  zero mode, rather than a  $\psi$  zero mode, we should focus on configurations that obey

$$\frac{dx^i}{d\tau} = g^{ij} \frac{\partial h}{\partial x^j} \quad (3.20)$$

That is, we should focus on instantons rather than anti-instantons. This means that we should look at configurations that start, at  $\tau \rightarrow -\infty$  at  $X_b$  and end up at  $\tau \rightarrow +\infty$  at  $X_a$ , with  $h(X_a) > h(X_b)$ . If we think of  $h(x)$  as a height function, these are trajectories that go up, rather than down.

### Instantons and the Relative Morse Index

We’ve now played the “Grassmann integration” card twice: once in (3.18) to argue that we should get contributions only between vacua that have relative Morse index 1, and again above to argue that we should only get contributions from instantons with  $\mathcal{I}(D^\dagger) = 1$ . Clearly we need these two different arguments to coincide. Happily they do because of the following result:

**Claim:** The index of  $D^\dagger$  is equal to the relative Morse index

$$\mathcal{I}(D^\dagger) = \mu(X_a) - \mu(X_b)$$


**Proof(ish):** Here we give a sketch of the proof of this statement. The operator  $D$ , defined in (3.19), acts on an  $n$ -dimensional space of fermions  $\psi^i$  and takes the form

$$D^\dagger = -\frac{d}{d\tau} + \text{Hess}[h]$$

where  $\text{Hess}[h]$  is the  $n \times n$  Hessian matrix

$$\text{Hess}[h]^i_j = g^{ik} \nabla_k \partial_j h$$

When evaluated at the critical points  $x = X_a$ , this coincides with the Hessian that we previously used to define the Morse index. But the equation above provides an



**Figure 11.** An example of the spectral flow of eigenvalues between two vacua, albeit one in which creative licence has trumped mathematical precision. The level crossings shown above can occur but they are not generic. In a more typical trajectory, the order of eigenvalues remains the same under spectral flow.

extension of the definition of the Hessian to each point along the instanton trajectory  $x(\tau)$ . As we move along this trajectory, the eigenvalues and orthonormal eigenvectors will smoothly evolve,

$$\text{Hess}[h(\tau)] \mathbf{e}_k(\tau) = \lambda_k(\tau) \mathbf{e}_k(\tau) \quad (3.21)$$

There is no sum over  $k = 1, \dots, n$  in this equation which labels the eigenvectors and eigenvalues. (The eigenvectors  $\mathbf{e}_k$  have an additional  $i = 1, \dots, n$  index which is the vector index and is suppressed in the equation above.)

We can now follow the  $n$  eigenvalues  $\lambda_i$  as we move from one critical point to another. This is known as *spectral flow*, and an example is shown in Figure 11. The number of negative eigenvalues at  $\tau = -\infty$  is the Morse index  $\mu(X_a)$ ; the number of negative eigenvalues at  $\mu = +\infty$  is  $\mu(X_b)$ .

Of particular interest are those eigenvalues which start negative and end up positive, or vice versa. The difference between those that cross in one direction, and those that cross in the other, is the relative Morse index  $\mu(X_a) - \mu(X_b)$ . The example shown in the Figure 11 has one more negative eigenvalue at the end than at the beginning which means that the corresponding instanton interpolates between two value with relative Morse index  $\mu(X_a) - \mu(X_b) = +1$ .

To solve the Dirac equation  $D^\dagger \psi = 0$ , we simply expand the fermions in terms of the eigenvectors, writing  $\psi^i(\tau) = \sum_k c_k(\tau) e_k^i(\tau)$ . We insert this ansatz into the Dirac equation and, using the orthogonality of eigenvectors  $g_{ij} e_k^i e_l^j = \delta_{kl}$ , we have

$$\left( \frac{d}{d\tau} - \lambda_k(\tau) \right) c_k(\tau) = - \sum_l g_{ij} e_k^i \dot{e}_l^j c_l \quad (3.22)$$

with no sum over  $k$ . First, let's suppose that we can ignore the term on the right-hand side. Then the equation has a straightforward solution,

$$c_k(\tau) = A_k \exp \left( + \int d\tau' \lambda_k(\tau') \right)$$

But this is a normalisable solution to the Dirac equation only if  $\lambda(\tau) < 0$  for  $\tau \rightarrow +\infty$  and  $\lambda(\tau) > 0$  for  $\tau \rightarrow -\infty$ . That is, we get a solution for every eigenvalue that flips from positive to negative. Meanwhile, the same analysis shows that every eigenvalue that goes the other way, from negative to positive, gives a solution to  $D\psi = 0$ . This is precisely what we wanted to show, namely

$$\mathcal{I}(D^\dagger) = \mu(X_a) - \mu(X_b)$$

That leaves us with the question of why it's legal to ignore the term on the right-hand side of (3.22). This is where we get to the “ish” part of proofish. The term captures how the eigenvectors twist as we move along the instanton trajectory due to the Levi-Civita connection. (In (3.22), this takes the form of a Berry connection.) This connection doesn't introduce any further topology into the game and it is a true fact that it doesn't change the index, albeit not a fact that I will demonstrate here. In acknowledgement of this slipshod approach, I'll replace the traditional QED box used at the end of a proof with something more wonky.  $\square$

It turns out that for background configurations with  $\mathcal{I}(D^\dagger) > 0$  we generically have  $\text{Ker } D = 0$ , so that  $\mathcal{I}(D^\dagger) = \dim \text{Ker } D^\dagger$ . (For example, the situation shown in Figure 11 is not generic.) We will assume that this is the case moving forwards.


The calculation above also tells us about the bosonic collective coordinates of an instanton. Suppose that we find a solution  $x(\tau)$  to the instanton equation (3.20). To see if this solution has any collective coordinates, we can look at variations  $x(\tau) + \delta x(\tau)$  and see if  $\delta x(\tau)$  satisfies the linearised instanton equation,

$$\frac{d}{d\tau} \delta x^i - g^{ij} \nabla_j \partial_k h \delta x^k = 0$$

But this coincides with the Dirac equation  $D^\dagger \delta \mathbf{x} = 0$  whose solutions we've just counted. The upshot is that the number of bosonic collective coordinates is equal to the number of fermi zero modes, and both are counted by the relative Morse index. A slicker way of saying this is to note that bosonic and fermionic zero modes are related by the unbroken supersymmetry  $Q^\dagger$  in the background of an instanton.

For our purposes, we want to consider instantons that interpolate between critical points with relative Morse index 1. Here the sole bosonic collective coordinate is the obvious one: the time  $\tau_1$  at which the instanton does its business of interpolating from one critical point to the other. This is the collective coordinate that we met previously in Section 2.2.

Although not of immediate utility, we can also get a feel for where the other bosonic collective coordinates may come from when  $\Delta\mu > 1$ . Consider the height Morse function on the round sphere  $\mathbf{S}^2$ . We know that there are two critical points at the south and north pole with Morse index 0 and 2 respectively. Correspondingly, the instanton that interpolates from the south to the north pole has two collective coordinates: one is the time  $\tau_1$  at which the instanton makes the jump, the other is the angle  $\phi$  of the trajectory as shown in the figure. In this example, the second collective coordinate is obvious because it arises due to a symmetry. But the arguments above tell us that, perhaps surprisingly, this second collective coordinate persists even when we deform the sphere, or potential, so that there's no longer a rotational symmetry.




### Completing the Instanton Computation

The rest of our instanton computation proceeds in exactly the same manner as that of Section 2.3. Our final answer is the obvious generalisation (2.41): for vacua  $|\Psi_a\rangle$  and  $|\Psi_b\rangle$ , whose Morse index differs by 1, we have

$$\langle \Psi_a | Q | \Psi_b \rangle = \frac{e^{-S_{\text{inst}}}}{\sqrt{2\pi}} \sum_{\gamma} n_{\gamma} \quad (3.23)$$

Here the sum is over all distinct instantons  $\gamma$  and  $n_{\gamma} = \pm 1$  is a sign that comes from computing the determinants

$$n_{\gamma} = \frac{\det D}{\sqrt{\det D^\dagger D}} = \pm 1$$



**Figure 12.** Two different instanton trajectories interpolating between  $X_b$  with  $\mu(X_b) = 0$  and  $X_a$  with  $\mu(X_a) = 1$ .

We met this sign before in Section 2.3 where we confessed that it is a little tricky to fix. Now it is time to make good on our promise of explaining where it comes from.

Let's start in the vacuum  $|\Psi_a\rangle$  localised at the end of the instanton trajectory at  $X_a$ . There are  $\mu(X_a)$  negative eigenvalues of the Hessian and their eigenvectors span a  $\mu(X_a)$ -dimensional space that we call  $V_a$ . The ground state  $|\Psi_a\rangle$  is associated to a  $\mu(X_a)$ -form and this induces an orientation on  $V_a$ .

The tangent to the instanton trajectory at  $X_a$  lies in the space  $V_a$  of negative eigenvectors. Let us call this tangent vector  $v$ . Generically,  $v$  will coincide with the eigenvector with largest negative eigenvalue  $\lambda_k$  (or smallest  $|\lambda_k|$ ) since this is usually the unique direction for which the eigenvalue flips sign by the time we reach  $X_b$  at the bottom. We denote the subspace of  $V_a$  that is orthogonal to  $v$  as  $\tilde{V}_a$ . There is a natural orientation on  $\tilde{V}_a$  that comes from taking the interior product  $\iota_v \Psi_a$ .

Now we propagate the space  $\tilde{V}_a$  along the instanton trajectory  $\gamma$ . We can do this, for example, by following the eigenvectors  $\mathbf{e}_k(\tau)$  of (3.21) corresponding to those eigenvalues that remain negative along the entire journey.

By the time we reach the end of the trajectory, the orientation on  $\tilde{V}_a$  that we started with gives an orientation on  $V_b$ , the space of negative eigenvectors of the Hessian at  $X_b$ . But there is a different way to define an orientation on  $V_b$ , which is that induced by the ground state  $|\Psi_b\rangle$  or, more precisely, the corresponding  $\mu(X_b)$ -form. The question is: do these two ways of defining an orientation coincide? If they do, we take  $n_\gamma = 1$ . If they do not, we take  $n_\gamma = -1$ .

For example, consider again the deformed bean-shaped sphere shown in Figure 12. There are two instanton trajectories that interpolate from the minimum  $X_b$  at the bottom to the saddle point  $X_a$  in the middle. At  $X_a$ , the tangent vectors to the two different instanton trajectories point in different directions, and that means that each instanton trajectory induces opposite orientations  $\iota_v \Psi_a$  on  $\tilde{V}_a$ . Correspondingly, one instanton will have  $n_\gamma = +1$  and the other  $n_\gamma = -1$ , and the two cancel out in (3.23). This same argument explains why the ground states are not lifted for the double well on a circle that we discussed in Section 2.3.4.

### 3.2.2 The Morse-Witten Complex

Let's recap. A Morse function gives us a collection of critical points. There are  $m_p$  critical points  $X$  with Morse index  $p = \mu(X)$  and, associated to each, there is an energy eigenstate  $|\Psi_a\rangle$  and an associated  $p$ -form. These can be thought of as a basis for an  $m_p$ -dimensional space that we will call  $C^p$ .

Not all the energy eigenstates  $|\Psi_a\rangle$  have vanishing energy or, equivalently, not all of them are annihilated by  $Q$ . But, as we saw in Section 2.3, they all have energy that is, at most, of order  $e^{-S_{\text{inst}}}$ . The tunnelling calculation that we've just done shows that,


$$\langle \Psi_a | Q | \Psi_b \rangle = \frac{e^{-S_{\text{inst}}}}{\sqrt{2\pi}} \sum_{\gamma} n_{\gamma} \quad \text{whenever } \mu(X_a) - \mu(X_b) = 1$$

If we think about  $Q$  as acting within this space of states, we can insert an “almost resolution” of the identity  $\mathbf{1} \approx \sum_a |\Psi_a\rangle \langle \Psi_a|$  to get

$$\begin{aligned} Q|\Psi_b\rangle &= \sum_a \langle \Psi_a | Q | \Psi_b \rangle |\Psi_a\rangle \\ &= \sum_{a: \mu_a = \mu_b + 1} \sum_{\gamma} \frac{n_{\gamma}}{\sqrt{2\pi}} e^{-(h(X_a) - h(X_b))} |\Psi_a\rangle \end{aligned}$$

Here the “almost resolution” of  $\mathbf{1}$  is because we've neglected all higher energy states. But their overlap with the low lying states  $|\Psi_a\rangle$  is exponentially suppressed and can be ignored. This means that we're left with an expression for the action of  $Q$  among the critical points. Neither the factor of  $\sqrt{2\pi}$ , nor the instanton action, are important for our present purposes and can be absorbed into the normalisation of the states. We then have

$$Q|\Psi_b\rangle = \sum_{a: \mu_a = \mu_b + 1} \sum_{\gamma} n_{\gamma} |\Psi_a\rangle$$



**Figure 13.** The four different instanton trajectories on the bean.

This is a map  $Q : C^p \rightarrow C^{p+1}$ . More abstractly, it can be viewed as a map between spaces of critical points. And, importantly, it satisfies  $Q^2 = 0$ . This means that we can define a chain complex (strictly a cochain complex), known as the *Morse-Witten complex*, or sometimes the *Morse-Smale-Witten complex*

$$0 \longrightarrow C^0 \xrightarrow{Q} C^1 \xrightarrow{Q} \dots \xrightarrow{Q} C^n \xrightarrow{Q} 0$$

The cohomology of  $Q$  describes the  $E = 0$  ground states of the system or, equivalently, the Betti numbers.

As an example we can look once again at the bean shaped manifold shown in Figure 13. We've already seen that the two instantons taking us from  $X_4$  to  $X_3$  cancel out, leaving us with

$$Q|\Psi_4\rangle = 0$$

This means that  $|\Psi_4\rangle$  is a true ground state of the system. There are also two instanton trajectories emanating from  $X_3$ , one to each of the peaks at  $X_1$  and  $X_2$ . These trajectories have different orientations, meaning

$$Q|\Psi_3\rangle = |\Psi_1\rangle - |\Psi_2\rangle$$

Finally, we have  $Q|\Psi_1\rangle = Q|\Psi_2\rangle = 0$  as both states are top forms. The true ground states lies in  $Q$ -cohomology and there are two of them: the 0-form  $|\Psi_0\rangle$  and the 2-form  $|\Psi_1\rangle + |\Psi_2\rangle$ . This reproduces the cohomology of  $\mathbf{S}^2$ .

### 3.3 The Atiyah-Singer Index Theorem

We now turn to a second application of supersymmetric quantum mechanics. We will study a version of supersymmetric quantum mechanics that yields the *Atiyah-Singer index theorem*. Before introducing the physics, we first explain what problem the index theorem addresses.

In  $n$  dimensions, where  $n$  is even, a Dirac spinor  $\chi$  has  $2^{n/2}$  components. In flat space, the free Dirac equation reads

$$\not{D}\chi = \gamma^a \partial_a \chi = 0 \quad (3.24)$$

Here the gamma matrices obey the usual Clifford algebra

$$\{\gamma^a, \gamma^b\} = 2\delta^{ab} \quad a, b = 1, \dots, n \quad (3.25)$$

The only solutions to (3.24) are constant spinors. That's a bit boring and, in  $\mathbb{R}^n$ , more than a bit non-normalisable. Things get more interesting when the fermion lives on a curved manifold  $M$ . To describe this situation, we first introduce vielbeins

$$g_{ij} e_a^i e_b^j = \delta_{ab}$$

The  $i, j$  indices are raised and lowered using the metric  $g_{ij}$  while the tangent space indices  $a, b$  are raised and lowered using  $\delta_{ab}$ . (See the lectures on [General Relativity](#) for more details.) The Dirac equation then takes the form

$$\not{D}\chi = \gamma^a e_a^i D_i \chi = 0 \quad (3.26)$$

with the covariant derivative given by

$$D_i = \partial_i + \frac{1}{2}(\omega_i)^{bc} S_{bc} \quad (3.27)$$

Here  $S_{ab}$  are the generators of  $SO(n)$  (strictly  $\text{Spin}(n)$ ) in the spinor representation

$$S_{ab} = \frac{1}{4}[\gamma_a, \gamma_b]$$

Meanwhile  $(\omega_i)^{ab}$  is the *spin connection*, defined by

$$(\omega_i)^a_b = e^a_j \partial_i e_b^j + \Gamma_{cb}^a e^c_i = e^a_j \nabla_i e_b^j$$

We can then ask: how many solutions there are to the Dirac equation (3.26)? This is where the Atiyah-Singer index theorem comes in. It relates the number of solutions to the Dirac equation to the topology of the underlying manifold. The purpose of this section is to give a physics derivation of the index theorem from supersymmetric quantum mechanics.

### 3.3.1 The $N = 1$ Sigma Model

Our quantum mechanics of choice has half the supersymmetry of the models that we've considered until now in this section. That is, we will have  $N = 1$  supersymmetry with a single real supercharge  $Q$ . We met some simple theories of this kind already in Section 1.4.3.

With  $N = 1$  supersymmetry, the sigma model action (3.6) is replaced by something that, at first glance, appears much simpler,

$$L = \int dt \frac{1}{2} g_{ij}(x) \dot{x}^i \dot{x}^j + \frac{i}{2} g_{ij} \psi^i \nabla_t \psi^j \quad (3.28)$$

The key difference is that the Grassmann variables are now Majorana modes

$$\psi^{\dagger i} = \psi^i$$

We no longer have the Riemann tensor interaction term, but the Levi-Civita connection still shows up, as before, in the kinetic term for the fermions,

$$\nabla_t \psi^i = \frac{d\psi^i}{dt} + \Gamma^i_{jk} \frac{dx^j}{dt} \psi^k \quad (3.29)$$

Although the action (3.28) has fewer interaction terms, it also has less symmetry. In particular, because the fermions are real we no longer have the  $U(1)$  symmetry that rotated the phase of the fermions. For our  $N = 2$  sigma model (3.6), this symmetry ensured that the energy eigenstates had a fixed number,  $p$  of excited fermions. Now that we no longer have this symmetry, we expect the energy eigenstates to involve a mixture of different fermions. The only protection we have comes from the  $(-1)^F$  symmetry that categorises states into  $\mathcal{H}_B$  with an even number of fermions and  $\mathcal{H}_F$  with an odd number.

We've already seen in Section 1.4.3 what emerges when we quantise the fermions so we will be brief here. The canonical anti-commutation relations are

$$\{\psi^i, \psi^j\} = g^{ij}$$

which is closely related to the Clifford algebra (3.25): the relationship between the fermions and gamma matrices involves a vielbein to accommodate the presence of the metric:  $\psi^i = e_a^i \gamma^a$ . This is telling us that, upon quantisation, the fermions will give  $2^{n/2}$  states which can be viewed as a Dirac spinor  $\chi$  living on the manifold  $M$ . While quantisation of the  $N = 2$  sigma model (3.6) gave us  $p$ -forms over the manifold, now we have a spinor.

The action (3.28) is invariant under the supersymmetry transformations

$$\delta x^i = \epsilon \psi^i \quad \text{and} \quad \delta \psi^i = i \epsilon \dot{x}^i$$

with the associated supercharge

$$Q = \frac{1}{2} g_{ij} \psi^i \dot{x}^j$$

The ground states of the quantum mechanics once again obey  $Q|\chi\rangle = 0$ . We can ask how this equation translates into the geometric language. The answer is clear: the fermion  $\psi^i$  in the supercharge is replaced by a gamma matrix, while the mechanical momentum  $\dot{x}$  is replaced by the appropriate covariant derivative, so that  $Q = i \not{D}$ . The upshot is that ground states of the quantum mechanics are given by solutions to the Dirac equation

$$\not{D}\chi = 0 \tag{3.30}$$

where the covariant derivative is (3.27), as appropriate for a spinor on a curved manifold  $M$ . The Hamiltonian is then  $H = Q^2 = -\not{D}^2$ .

We can now see why this quantum mechanics is of interest. The ground states are specified by solutions to the Dirac equation which is exactly what we want to count. Moreover, we know how to count ground states in supersymmetric quantum mechanics, at least up to sign: we use the Witten index.

To get an expression for the index, we first need to figure out which states sit in  $\mathcal{H}_B$  and which in  $\mathcal{H}_F$ . As we already saw in Section 1.4.3, this has a particularly nice interpretation in terms of the spinor. Because we are an even dimension  $n$ , the Dirac spinor decomposes into two Weyl spinors which are eigenspinors of

$$\hat{\gamma} = i^{n/2} \gamma^1 \dots \gamma^n$$

This obeys  $\hat{\gamma}^2 = 1$  and  $\{\hat{\gamma}, \gamma^i\} = 0$ , and is the generalisation of the “ $\gamma^5$ ” matrix in four dimensions. This is the operator that determines whether a given state lies in  $\mathcal{H}_B$  or  $\mathcal{H}_F$  via the identification

$$\hat{\gamma} = (-1)^F$$

We can always pick a basis of gamma matrices that are block off-diagonal, so that we have

$$\hat{\gamma} = \begin{pmatrix} \mathbf{1} & 0 \\ 0 & -\mathbf{1} \end{pmatrix}$$

where each entry is a  $2^{n/2-1}$  dimensional matrix. The Dirac operator then takes the form

$$\not{D} = \begin{pmatrix} 0 & \mathcal{D}^\dagger \\ \mathcal{D} & 0 \end{pmatrix} \quad (3.31)$$

where  $\mathcal{D} : \mathcal{H}_B \rightarrow \mathcal{H}_F$  and  $\mathcal{D}^\dagger : \mathcal{H}_F \rightarrow \mathcal{H}_B$ . The Witten index coincides with the index of the operator  $\mathcal{D}$ ,

$$\text{Tr} (-1)^F e^{-\beta H} = \mathcal{I}(\mathcal{D}) := \dim \text{Ker } \mathcal{D} - \dim \text{Ker } \mathcal{D}^\dagger \quad (3.32)$$

This is also the quantity of relevance to the Atiyah-Singer index theorem. Our next task is to compute it. But, by now, our strategy for this should be clear: we turn to the path integral.

### 3.3.2 The Path Integral Again

The same argument that we used in Section 3.1.3 when deriving the Chern-Gauss-Bonnet theorem tells us that the path integral localises on constant configurations  $\dot{x}^i = \dot{\psi}^i = 0$ . We'll pick a constant configuration and expand

$$x^i(\tau) = x_0^i + \delta x^i(\tau) \quad \text{and} \quad \psi^i(\tau) = \psi_0^i + \delta \psi^i(\tau)$$

We then compute the path integral by performing a Gaussian integration over the fluctuations  $\delta x$  and  $\delta \psi$ . Our life is made easier if we work in normal coordinates in which

$$g_{ij}(x) = \delta_{ij} - \frac{1}{3} R_{iklj}(x_0) \delta x^k \delta x^l$$

and

$$\delta \Gamma_{jk}^i(x) = \partial_l \Gamma_{jk}^i(x_0) \delta x^l = -\frac{1}{3} (R^i{}_{jkl}(x_0) + R^i{}_{kjl}(x_0)) \delta x^l$$

To quadratic order, the Euclidean action then becomes

$$S_E = \frac{1}{2} \int d\tau \delta_{ij} \left( -\delta x^i \frac{d^2}{d\tau^2} \delta x^j + \delta \psi^i \frac{d}{d\tau} \psi^j \right) - \frac{1}{2} R_{ijkl} \psi_0^i \psi_0^j \delta x^k \frac{d\delta x^l}{d\tau}$$

Performing the Gaussian integral periodic boundary conditions, we have

$$Z = \int \mathcal{D}\delta x \mathcal{D}\delta \psi e^{-S_E} = \sqrt{\frac{\det'(\delta_{ij} \partial_\tau)}{\det'(-\delta_{ij} \partial_\tau^2 + \Omega_{ij} \partial_\tau)}} = \sqrt{\frac{1}{\det'(-\delta_j^i \partial_\tau + \Omega_j^i)}}$$

The fermionic determinant in the numerator now sits under square root, reflecting the fact that the fermions are real. (It could be better thought of as a Pfaffian.) Both determinants have had their zero modes truncated since these correspond to the integrals over  $x_0$  and  $\psi_0$ , both of which we will do explicitly below.

After the small cancellation seen above, we're left with the task of computing the determinant operator involving the matrix

$$\Omega_j^i := \delta^{ip} R_{pjkl}(x_0) \psi_0^k \psi_0^l \quad (3.33)$$

This should be thought of as an  $n \times n$  matrix that depends both on the point  $x_0$  and on the background fermion  $\psi_0$ . (It may seem odd to think about a matrix as depending on a background Grassmann parameter like  $\psi_0$ ; the meaning of this should become clearer below when we think about what we're going to do with this matrix.) This is an anti-symmetric matrix and we can choose a basis in which it takes block diagonal form

$$\Omega_j^i = \begin{pmatrix} \mathbb{W}_1 & & & \\ & \mathbb{W}_2 & & \\ & & \ddots & \\ & & & \mathbb{W}_{n/2} \end{pmatrix} \quad \text{with } \mathbb{W}_a = \begin{pmatrix} 0 & \omega_a \\ -\omega_a & 0 \end{pmatrix}$$

The eigenvalues are  $\pm i\omega_a$  and these depend on both  $x_0$  and on  $\psi_0$ . We can also diagonalise the derivative term simply by working in a Fourier basis around the circle. Since we know that the end result for the Witten index must be independent of  $\beta$ , we'll take advantage of this and work with  $\beta = 1$ . We then have

$$\delta x^i(\tau) \sim e^{ik\tau} \quad \text{with } k = 2\pi p \quad \text{and } p \in \mathbb{Z}$$

This means that the eigenvalues of the bosonic fluctuation operator are  $i(k \pm \omega)$ . Restricting to any given  $2 \times 2$  matrix  $\mathbb{W}$ , we have

$$\begin{aligned} \sqrt{\det'(-\partial_\tau + \mathbb{W})} &= \prod_{p \neq 0} (2\pi ip + i\omega)^{1/2} (2\pi ip - i\omega)^{1/2} \\ &= \prod_{p=1}^{\infty} (2\pi ip)^2 \left[ 1 + \left( \frac{i\omega}{2\pi p} \right)^2 \right] \end{aligned}$$

We've met each of these products before. The second, convergent product is given by (2.6)

$$\prod_{p=1}^{\infty} \left[ 1 + \left( \frac{i\omega}{2\pi p} \right)^2 \right] = \frac{\sinh(i\omega/2)}{i\omega/2}$$

The first, divergent, product can be treated using zeta function regularisation as in (2.7) and gives

$$\prod_{p=1}^{\infty} (2\pi ip)^2 = -i$$

Putting this together, we have an expression for the partition function after integrating over the fluctuations,

$$Z = (-i)^{n/2} \prod_{a=1}^{n/2} \frac{i\omega_a/2}{\sinh(i\omega_a/2)}$$

We're now left just with the zero mode integrations: including these gives us the expression for the Witten index and hence the index of the Dirac operator

$$\mathcal{I}(\mathcal{D}) = (-i)^{n/2} \int \prod_{i=1}^n \frac{dx_0^i}{\sqrt{2\pi}} d\psi_0^i \prod_{a=1}^{n/2} \frac{i\omega_a/2}{\sinh(i\omega_a/2)}$$

The next step is to do the fermion zero mode integration. The idea here is that each  $\omega_a$  depends quadratically on the fermion zero modes. We should expand each term in the product,

$$\frac{\omega/2}{\sinh(\omega/2)} = 1 - \frac{\omega^2}{24} + \frac{7\omega^4}{5760} + \dots$$

Because the function is even, the expansion contains only even powers of  $\omega$  and hence the fermionic variables  $\psi_0$  come in groups of four. The fermionic integration picks out the term that saturates the Grassmann integration. We learn that the index  $\mathcal{I}(\mathcal{D})$  will be non-vanishing only on manifolds whose dimension  $n$  is a multiple of four. (This also eliminates the factors of  $i$  in the expression for  $\mathcal{I}(\mathcal{D})$ .)

There is a more geometric way to think about this. Instead working with fermions, we turn again to forms. (The fact that our fermions don't yield forms upon quantisation is irrelevant here: this is just a formal trick.) We introduce the curvature 2-form

$$\mathcal{R}^a_b = R^a_{bcd} \hat{\theta}^c \wedge \hat{\theta}^d$$

where  $\hat{\theta} = e^a_i dx^i$  are a basis of one-forms. (see the lectures on [General Relativity](#) for more details.) Clearly  $\mathcal{R}$  has the same formal structure as the fermionic matrix  $\Omega$  that we met before. This means that we can equally well write

$$\mathcal{I}(\mathcal{D}) = \int_M \hat{A}(M) \quad \text{with} \quad \hat{A}(M) = \frac{1}{(2\pi)^{n/2}} \sqrt{\det \left( \frac{\mathcal{R}/2}{\sinh(\mathcal{R}/2)} \right)}$$

This is the Atiyah-Singer index theorem. The expression  $\hat{A}(M)$  is referred to as the *A-roof genus* or *A-hat genus* of  $M$  (or, more correctly, the tangent bundle of  $M$ ). The expression should be viewed as expanding out the determinant until we find a top form

that can be integrated over the manifold  $M$ . The terms that appear in this expansion are

$$\int_M \hat{A}(M) = 1 - \frac{p_1}{24} + \frac{1}{16} \left( \frac{7}{360} p_1^2 - \frac{1}{90} p_2 \right) + \dots$$

where the various terms in the expansion arise for manifolds of dimension  $n = 0, 4, 8$  and so on. The  $p_i$  are known as *Pontryagin numbers* and can be expressed as integrals of the curvature 2-form over the manifold  $M$ . For a manifold of dimension  $\dim(M) = 4$ , we have

$$\int_M \hat{A}(M) = -\frac{p_1}{24} \quad \text{with} \quad p_1 = -\frac{1}{8\pi^2} \int_M \text{tr} \mathcal{R}^2 \quad (3.34)$$

The simplest examples of 4-manifolds are the torus and sphere, which have  $p_1(\mathbf{T}^4) = p_1(\mathbf{S}^4) = 0$ . This tells us that the index of the Dirac operator vanishes. For the torus  $\mathbf{T}^4$ , with periodic boundary conditions for spinors, this is because both  $\mathcal{D}$  and  $\mathcal{D}^\dagger$  have zero modes. For the sphere  $\mathbf{S}^4$ , it is because there are no zero modes.

To find a compact manifold  $M$  with  $p_1 \neq 0$ , we need to turn to something more exotic. A nice example is provided by the manifold known as  $M = K3$ , which can be viewed as a smooth quartic surface in  $\mathbf{CP}^3$ . This is the only non-trivial Calabi-Yau 4-manifold and has  $p_1(K3) = -48$ .

The factor of  $1/24$  in (3.34) is telling us something interesting. Because the left-hand side is counting something, the right-hand-side must be an integer. This suggests that the integral  $p_1$  must be a multiple of 24. In fact, things are a little more subtle. The thing that we're counting is the number of solutions to the Dirac equation and to pose this question at all, we must be able to put Dirac spinors on the manifold  $M$ . This, it turns out, is not possible for all manifolds. Those for which it is possible are called *spin manifolds*. And for any orientable spin manifold, it turns out that  $p_1$  is always divisible by 48.

The canonical example of a non-spin 4-manifold is complex projective space  $\mathbf{CP}^2$ . It's not possible to consistently patch spinor fields over this space, and so the question of counting solutions to the Dirac equation is irrelevant. It turns out that  $p_1(\mathbf{CP}^2) = 3$ . Moreover, one can show that for any orientable 4-manifold,  $p_1$  is always divisible by 3.

For manifolds with  $\dim M = 8$ , we need the result

$$p_2 = \frac{1}{128\pi^4} \int_M \left( (\text{tr} \mathcal{R}^2)^2 - 2\text{tr} \mathcal{R}^4 \right)$$

There are generalisations to higher dimensional manifolds.

### 3.3.3 Adding a Gauge Field

There is an interesting generalisation of the action (3.28) that retains  $N = 1$  supersymmetry. This is

$$L = \int dt \frac{1}{2} g_{ij}(x) \dot{x}^i \dot{x}^j + \frac{i}{2} g_{ij} \psi^i \nabla_t \psi^j + i \eta_\alpha^\dagger D_t \eta^\alpha + \frac{1}{2} (F_{ij})^\alpha{}_\beta \eta_\alpha^\dagger \eta^\beta \psi^i \psi^j \quad (3.35)$$

with  $i, j = 1, \dots, n$  as before and  $\alpha, \beta = 1, \dots, r$ . Here the  $\psi^i$  are Majorana fermions, with covariant derivative given by the same expression (3.29) that we had before. Meanwhile, the  $\eta_\alpha$  are complex fermions with covariant derivative

$$D_t \eta^\alpha = \frac{d\eta^\alpha}{dt} + (A_i)^\alpha{}_\beta \frac{dx^i}{dt} \eta^\beta$$

We should think of  $A_i$  as a  $U(r)$  gauge connection over the manifold  $M$ . Just as the metric  $g_{ij}$  is something fixed, so too is this gauge field. In more mathematical language, we should think of  $A$  as a connection of a vector bundle  $E$ . In the action (3.35)  $F$  is the associated field strength

$$(F_{ij})^\alpha{}_\beta = \partial_i (A_j)^\alpha{}_\beta - \partial_j (A_i)^\alpha{}_\beta + [A_i, A_j]^\alpha{}_\beta$$

Note that the four-fermion term involves  $F$ , which is the curvature of  $A$ . This is entirely analogous to the situation in  $N = 2$  supersymmetric quantum mechanics where the four-fermion term involves the Riemann tensor, which is the curvature of the spin connection.

At first glance, it is surprising that the action (3.35) admits supersymmetry. Until now, all our examples of supersymmetry involve a matching between bosonic and fermionic degrees of freedom. But here we have introduced additional fermionic degrees of freedom  $\eta^\alpha$  without the corresponding bosonic partners. Nonetheless, you can check that the action is invariant under the following  $N = 1$  supersymmetry,

$$\begin{aligned} \delta x^i &= \epsilon \psi^i \\ \delta \psi^i &= i \epsilon \dot{x}^i \\ \delta \eta^\alpha &= -\epsilon \psi^i (A_i)^\alpha{}_\beta \psi^\beta \end{aligned}$$

Together with the conjugate expression  $\delta \eta_\beta^\dagger = -\epsilon \eta_\alpha^\dagger (A_i)^\alpha{}_\beta \psi^i$ . It turns out that adding extra fermions, without bosonic counterparts, is an option only for supersymmetric theories in  $d = 0 + 1$  and  $d = 1 + 1$  dimensions.

How should we think about the theory (3.35). Quantising the real fermions  $\psi^i$  gives us a spinor over  $M$  as before. Quantising the complex fermions  $\eta^\alpha$  gives us forms, but they're not forms over the manifold  $M$  since they don't carry the  $i = 1, \dots, n$  index. Instead, they are forms over the gauge bundle  $E$ . In more mundane language, this simply means that the states transform in different representations of the  $U(r)$  gauge connection. Ignoring the  $\psi^i$  fermions for now, we start with the state  $|0\rangle$  such that

$$\eta^\alpha|0\rangle = 0$$

This is a singlet (i.e. neutral) under the  $U(r)$  gauge bundle. Next, we have  $\eta^{\dagger\alpha}|0\rangle$  which sit in the fundamental representation of  $U(r)$ , and the  $\eta^{\dagger\alpha}\eta^{\dagger\beta}|0\rangle$  which sits in the anti-symmetric representation, and so on. Furthermore, there is a  $U(1)$  symmetry that rotates  $\eta^\alpha \rightarrow e^{i\theta}\eta^\alpha$  and this ensures that energy eigenstates have fixed number of  $\eta^\dagger$  excitations and so sit in a fixed representation of  $U(r)$ .

The upshot is that the Hilbert space consists of a collection of spinors over  $M$ , each transforming in the  $p^{\text{th}}$  antisymmetric representation of  $U(r)$ . The  $E = 0$  ground states obey  $Q|\chi\rangle = 0$  which, translated into the geometric language, becomes the Dirac equation  $\not{D}\chi = 0$  with

$$D_i = \partial_i + \frac{1}{2}(\omega_i)^{bc}S_{bc} + A_i$$

with  $A_i$  in the appropriate representation. Once again, this can be put in block off-diagonal form (3.31) and we can compute the index  $\mathcal{I}(\mathcal{D})$  which, as in (3.32), coincides with the Witten index. Restricting to the fundamental representation, a similar calculation to the one above now yields the Atiyah-Singer index theorem

$$\mathcal{I}(\mathcal{D}) = \int_M \hat{A}(M) \wedge \text{ch}(F) \tag{3.36}$$

where  $F$  is the *Chern character*,

$$\text{ch}(F) = \text{Tr} e^{F/2\pi} = r + c_1(F) + c_2(F) + \dots$$

The individual Chern classes are topological invariants of the gauge field when integrated over the manifold  $M$ . The first two are

$$c_1 = \frac{1}{2\pi} \text{Tr} F \quad \text{and} \quad c_2 = \frac{1}{8\pi^2} \text{Tr} F \wedge F - \text{Tr} F \wedge \text{Tr} F$$

There are close connections here to the physics of solitons. In particular, the integral of the first Chern number counts the number of vortices in a gauge theory. The integral of the second Chern number counts the number of Yang-Mills instantons. Both of these have an interesting zero mode structure, even in flat spacetime, as follows from the Atiyah-Singer index theorem (3.36).

### 3.4 What Comes Next?

We have barely scratched the surface of supersymmetric theories and their connection to mathematics. In this final section we give a brief sketch of the next steps. Roughly speaking, there are two directions that are particularly rich: increase the number of supersymmetries, and increase the spacetime dimension of the quantum theory.

First the supersymmetries. The stories we told above revolved around  $N = 2$  supersymmetry (for Morse theory) and  $N = 1$  supersymmetry (for the index theorem). We briefly met theories with  $N = 4$  supersymmetry in Section 1.4.2 where we saw that they naturally come with complex fields and a holomorphic superpotential. This suggests that a sigma model with  $N = 4$  supersymmetry should have a target space  $M$  that is, in some sense, a complex manifold.

This is indeed what happens. Sigma models with  $N = 4$  supersymmetry have target spaces that are *Kähler*. These target spaces necessarily have even dimension, with coordinates that can be paired together into complex numbers consistently over the entire manifold. This structure is best seen through the introduction of superfields, where the kinetic terms in the Lagrangian are written directly in terms of a function known as the *Kähler potential*  $K(\phi, \bar{\phi})$  that is related to the metric by

$$g_{i\bar{j}} = 2 \frac{\partial^2 K}{\partial \phi^i \partial \bar{\phi}^{\bar{j}}} \quad (3.37)$$

Superfields and the Kähler potential are both described in the accompanying lectures on [Supersymmetric Field Theory](#).

More supersymmetry brings yet more structure to the geometry, at least to a point. There are theories with  $N = 8$  supersymmetry whose target spaces are *hyperKähler*. Such manifolds have a dimension that is a multiple of four, with three inequivalent ways of pairing coordinates together into complex numbers. It's a little like having a quaternionic structure on the manifold (although beware that there is a different kind of object in mathematics known as a “quaternionic manifold”). However, the interesting things don't keep happening forever and by the time we get to  $N = 16$  supersymmetry the restriction becomes too strong and the target space is obliged to be flat and largely boring.

The full riches of supersymmetry really come when we consider theories in higher dimensions, meaning quantum field theories rather than quantum mechanics. While there are interesting stories for field theories in any spacetime dimension  $d \leq 6$  (and, if you include gravitational theories, for  $d \leq 11$ ) there is, as I now explain, a reason why QFTs in  $d = 1 + 1$  dimensions are special.

Consider a sigma model in  $d$  spacetime dimensions. We introduce coordinates  $x^\mu$ , with  $\mu = 0, 1, \dots, d-1$  for spacetime. The action is again based on some target space  $M$ . This means that the fields  $\phi^i(x)$ , with  $i = 1, \dots, n$  should be thought of as coordinates on  $M$ . The sigma model takes the form

$$S = \int d^d x g_{ij}(\phi) \partial_\mu \phi^i \partial^\mu \phi^j + \text{fermions} \quad (3.38)$$

Note that there are two conceptually different spaces in this action. The spacetime of the quantum field theory has dimension  $d$ , while the target space  $M$  has dimension  $\dim M = n$ .

With no potential  $V(\phi)$  to preference one point on the manifold  $M$  from another, this theory has many classical ground states: each point on  $M$  or, equivalently, each constant value of  $\phi^i$  should be viewed as a different classical ground state of the system.

But what happens in the quantum theory? We've already seen that for  $d = 1$ , which is just quantum mechanics, the ground state wavefunction spreads over  $M$ . This is important as it means that the ground state knows something about the entire manifold  $M$  and may therefore encode some information about its topology. We've seen examples of this throughout these lectures.

What happens in higher dimensions with  $d > 1$ ? It turns out that  $d = 2$  is just like quantum mechanics: the ground state wavefunction spreads over the whole manifold  $M$ . Meanwhile, at least in this respect, quantum theories in dimensions  $d \geq 3$  behave like the classical theory: each point on  $M$  defines a different ground state.

I won't prove this statement in these lectures. It sometimes goes by the name of the (Coleman)-Mermin-Wagner theorem and is closely related to the concept of a "lower critical dimension" in [Statistical Field Theory](#). At heart, it boils down to a property of the Poisson equation  $\nabla^2 \phi = \delta(x)$  in  $d$  Euclidean dimensions. At long distances, the solution grows in  $d = 1$  and  $d = 2$ . (It is  $\phi \sim x$  in  $d = 1$  and  $\phi \sim \log x$  in  $d = 2$ .) Conversely, the solution decays at long distance as  $\phi \sim 1/x^{d-2}$  in  $d \geq 3$ . Physically, this translates into the fact that wavefunctions spread over  $M$  for sigma models with  $d = 1$  and  $d = 2$ , while the ground state remains localised at a point in  $M$  for  $d \geq 3$ .

This means that if we want to find some interesting physics that captures topological properties of  $M$ , we should first look at  $d = 1$  and  $d = 2$ . We've now spent over 100 pages studying supersymmetric quantum mechanics. So the next step is to look at supersymmetric sigma models of the form (3.38) in  $d = 1 + 1$ .

There is a rather special feature of quantum field theories in  $d \geq 2$  dimensions that distinguishes them from quantum mechanics in  $d = 1$ : renormalisation. This means that the coupling constants that characterise a theory are not, in fact, constant. Instead they change with scale. Sigma models in  $d = 1 + 1$  dimensions are no exception. Here, the couplings of the theory are encoded in the metric  $g_{ij}(\phi)$ . Under renormalisation, this metric changes and depends on the scale  $\mu$  at which you look. The manner in which the metric changes is governed by the beautifully geometric beta function equation, known as *Ricci flow*

$$\mu \frac{\partial g_{ij}}{\partial \mu} = R_{ij} \tag{3.39}$$

This equation also plays an important role in [String Theory](#), where it is ultimately responsible for the emergence of the Einstein equations of general relativity.

Taking into account the renormalisation (3.39), there are three types of behaviour that can occur depending on the type of target space  $M$ . Those spaces with positive Ricci curvature,  $R > 0$ , will shrink under RG flow. In this case, the theory becomes increasingly strongly coupled in the infra-red and the impact on the physics is rather dramatic with the seemingly massless scalars  $\phi^i$  developing a quantum-generated mass. Examples include  $M = \mathbf{S}^n$  and  $M = \mathbf{CP}^n$ . You can read more about this in the lectures on [Statistical Field Theory](#) and the lectures on [Gauge Theory](#).

Target spaces  $M$  with negative Ricci curvature,  $R < 0$ , will typically expand under RG flow. In this case they become more and more weakly coupled as the flow to they infra-red. Hyperbolic spaces provide a simple example.

The sweet spot are those target spaces  $M$  for which the metric is Ricci flat, with  $R_{ij} = 0$ . In fact, rather wonderfully, you don't have to start in the UV with a manifold  $M$  with a Ricci flat metric. If the manifold admits a Ricci flat metric, then the quantum theory will typically find it through the RG flow. The long-wavelength physics of such a theory is then governed by an interacting conformal field theory. Or, if we're dealing with a supersymmetric sigma model, an interacting supersymmetric conformal field theory or SCFT for short.

At this point, there is again an lovely intersection with results from mathematics. If we have  $N = 4$  supersymmetry so that the target space is Kähler, then there is famous class of compact manifolds  $M$  that admit a Ricci flat metric known as *Calabi-Yau manifolds*. (For what it's worth, they are defined by having vanishing first Chern class.) Our discussion above means that for each Calabi-Yau manifold  $M$  there is an associated SCFT.

That, it turns out, is interesting. While the quantum mechanical sigma models described earlier in these lecture notes capture well-known stories of geometry, the  $d = 1 + 1$  sigma models give a new lens through which to look at the manifolds  $M$ . And this lens gives us new information about the manifolds that goes beyond what mathematicians originally knew (although they very quickly caught up!) It is appropriate to refer to this view of the target space  $M$ , as seen by a SCFT, as “quantum geometry”.

There are many novelties that come from associating a SCFT to a Calabi-Yau manifold  $M$ . But one stands out. There is not a unique association between manifolds  $M$  and SCFTs. Instead, pairs of manifolds  $M$  and  $N$  turn out to give rise to the *same* SCFT. These two manifolds  $M$  and  $N$  are topologically distinct and, naively, one wouldn't have thought that they have anything to do with each other. But, perhaps surprisingly, “quantum geometry” turns out to be more myopic than classical geometry and the SCFT approach cannot distinguish objects which appear obviously different to a classical geometer. At first glance, this myopia might appear to be a weakness, but closer examination shows that it is very much a strength. The myopia only arises because there are deep connections between the two manifolds  $M$  and  $N$ , with the geometric information of one encoded in a hidden and subtle form in the other. In technical language, the complex structure of one manifold is mapped to the symplectic structure of the other. This mapping is known as *mirror symmetry*. It is, sadly, a topic for another course.